

UC-NRLF



B 4 251 244

THE  
STAR - GUIDE

LATIMER CLARK F.R.A.S.

AND

HERBERT SADLER F.R.A.S.

*New*

UNIVERSITY OF CALIFORNIA.

FROM THE LIBRARY OF  
WILLIAM M. PIERSON.

GIFT OF MRS. PIERSON AND L. H. PIERSON.

No.

*Wm M Pierson*

*Dec 1/89*

SAMUEL GARDNER  
PUBLISHER  
WHOLESALE BOOKSELLER  
208 Post St





THE  
STAR-GUIDE

311



THE  
STAR-GUIDE

A LIST OF THE  
MOST REMARKABLE CELESTIAL OBJECTS  
VISIBLE WITH SMALL TELESCOPES  
WITH  
THEIR POSITIONS  
FOR EVERY TENTH DAY IN THE YEAR  
AND  
OTHER ASTRONOMICAL INFORMATION

BY  
LATIMER CLARK, F.R.A.S.  
AND  
HERBERT SADLER, F.R.A.S.



LONDON  
MACMILLAN AND CO.

1886

Q36A  
C6

PRINTED BY  
SPOTTISWOODE AND CO., NEW-STREET SQUARE  
LONDON



## INTRODUCTION.

---

THE first portion of this little work is intended primarily to serve as an introduction to the sidereal portion of 'Celestial Objects for Common Telescopes,' the well-known work of the late deeply regretted Prebendary Webb, and is designed for the use of those who possess small telescopes of from two to four inches aperture and upwards, by reminding them what objects to look for at each period of the year, and where to look for them. Many amateurs who are just commencing the study of Astronomy experience some difficulty in the selection of those objects, out of the countless number in the heavens, which are best suited for observation with small telescopes, and are apt to strain their eyes and waste their time in fruitless endeavours to catch faint nebulae or to separate difficult double stars, which can only really be observed with much larger telescopes than most beginners usually possess. It is hoped, therefore, that to such, a carefully selected list, on a definite plan, excluding all objects which are beyond the grasp of small telescopes, and including nearly all those which can be profitably examined with instruments of from two to three inches of aperture, may be of use. With this view a list of nearly six hundred of the most interesting and beautiful objects visible in the northern heavens with instruments of this description, has been arranged in a tabular form in order of Right Ascension, which the authors hope will be found to contain nearly every double star between the North Pole and  $15^{\circ}$  South Declination, which can be seen easily with an achromatic of small aperture. The positions and times of transit are given for each tenth day in the year. With very few exceptions no double star

has been inserted the brightest component of which falls below the seventh magnitude—the magnitude of the smallest of the satellites of Jupiter—or the fainter of which is not brighter than a tenth magnitude star, or any pairs which are too close to be conveniently seen in a telescope of small size. No red star has been included the colour of which cannot be seen, or any nebula or cluster which cannot conveniently be observed with such an instrument.

In the first column of the following list of objects visible with small telescopes the name of the star is given; all those to which Bayer assigned a Greek letter, or those numbered and catalogued by Flamsteed, Bode, or Piazzi, being designated by such letter or number. Flamsteed's numbers precede the name of the constellation to which they refer, *e.g.* 77 Pegasi; but as all Bayer's lettered stars have also numbers assigned to them by Flamsteed, the latter have been omitted. Piazzi's are numbered according to the hour of Right Ascension in which they happened to fall in the year 1800, *e.g.* P. iii. 98 (Eridani); while Bode's numbers follow the name of the constellation, as Coronæ 1. In one or two cases the star has been entered according to its number in the catalogue published by the British Association, as B. A. C. 352 (Andromedæ). In order to avoid all unnecessary technicalities, where a star, either from its comparative faintness or for some other reason, has failed to receive a Greek letter or one of Flamsteed's or Piazzi's numbers, it has been entered simply as a red star or pair in such and such a constellation. Thus the seventh object in the list is entered as 'Pair in Cassiopeia,' with a Right Ascension of  $23^{\text{h}} 55^{\text{m}} 5^{\text{s}}$ , and a North Declination of  $59^{\circ} 43'$ . The principal star of this pair is known to astronomers as A.Oe<sub>1</sub>. 26,287 (or O. Arg. N. 26,287), meaning that it is number 26,287 in a catalogue of stars between  $45^{\circ}$  and  $80^{\circ}$  North Declination, whose places were determined by Argelander, and that the catalogue was edited by Oeltzen, in the third series of the Annals of the Vienna Observatory, from which we have deduced its apparent place in the list for January 1886. As a *pair* it is termed OΣΣ (or OΣ<sup>2</sup>) 254, which refers to its number, 254, in a catalogue of double stars observed by O. Struve. Such details, however, are out of place in a list like the present. In the same manner nebulae

and clusters have been entered as 'Cluster in Cepheus,' 'Nebula in Cetus,' &c., without giving their numbers in the catalogues of Messier, Sir W., or Sir J. Herschel.

The second column is headed 'Visible,' and is simply intended to show observers at a glance during what months of the year the several objects in the lists may be most conveniently observed.

The third and fourth columns contain the Right Ascension and Declination of the objects, given to the nearest tenth of a minute of time in Right Ascension and the nearest minute of arc in Declination, and reduced to January 1, 1886. All the star places throughout the book have been brought up to this epoch.

The next four columns contain the sidereal time at 9 P.M. on the 1st, 11th, 21st, and last day of every month, and the mean time of transit at Greenwich of every object on those days. Thus the list for each month includes nearly fifty objects which pass the meridian at convenient hours for observation. The observer, however, is by no means confined to the list for each month alone, or to the southern part of the heavens. He may often desire to examine objects in the eastern skies, which may be found, not in the list for the month, but in the lists for the next following month; or he may desire to observe western stars, which will be found in the preceding months. These lists should, therefore, constantly be referred to. The stars transit every day  $3^m 55^s 91^s$  (or say 4 minutes) earlier than on the preceding day. A star which souths at ten to-night must, therefore, have southed two hours later a month ago, and will south two hours earlier a month hence. Remembering this, and looking at the position of the star, we can easily judge which list to consult. Suppose that on October 20 we desire to observe certain stars in the western heavens. We know from their position that they will be found in the list of stars some two or three months before; we therefore search the July and August lists, and adding, say, 31 days for August, 30 for September, and 20 for October, we find the interval since July 31, to be 81 days. Allowing  $3^m 56^s$  per diem for acceleration,<sup>1</sup> we get  $5^h 18^m$  to be *subtracted* from the times given on July 31, and

<sup>1</sup> For the convenience of reckoning a table of these accelerations will be found at page 26.

this will give us the time of southing of all the stars in the July list which are still visible to us. Similarly, for eastern skies, we know that the stars will south at convenient hours some two or three months hence, and will be found probably in the January list. We count, therefore, 72 days to January 1, and *adding*  $4^h 43^m$  to any of the stars in that list, we get their time of southing this evening. Before observing it is convenient to take out these times from the table, and to write them (in an abbreviated form) on a card. Thus for October 20 we have—


List of Stars	Interval in Days	Time to be SUBTRACTED	List of Stars	Interval in Days	Time to be ADDED
July 31 . . .	81	h. m. 5 18	November 1 .	11	h. m. 0 43
August 31 . .	50	3 17	December 1 .	41	2 41
September 30	20	1 19	January 1 . .	72	4 43

We are now in a position to find instantly the time of transit of any object in any of the lists from July to January. In making these notes the observer of course includes at the same time the correction (if necessary) due to his difference in longitude. If his station be westward of Greenwich, he must subtract the correction from the left hand column and add it to the right, and *vice versa*.

The next, or ninth, column contains the distances of the double stars, measured from centre to centre of the stars, and expressed in seconds and tenths of a second of arc. The succeeding column contains the position angle, or the angle of inclination to the meridian, of the smaller star as referred to the larger one of the pair. The following diagram, in which the arrow indicates the direction of motion through the field, will serve to explain the method of expressing position angles at present adopted. Up to about 1820, it was usual to reckon by quadrants, *i.e.*  $10^{\circ}nf$  answered to  $80^{\circ}$  in our present mode of reckoning,  $10^{\circ}sf$  to  $100^{\circ}$ ,  $10^{\circ}sp$  to  $260^{\circ}$ , and  $10^{\circ}np$  to  $280^{\circ}$ . It is, perhaps, as well to remind observers that the present mode of reckoning position angles is not that used in the prediction of occultations in the 'Nautical Almanac,'

<sup>1</sup> For the convenience of reckoning a table of these accelerations will be found at page 26.

where the initial point is taken at the top, or south, and the angles are reckoned round towards the right hand in the direction of the



revolution of the hands of a watch,  $180^\circ$  being at the north, or bottom. Great care has been taken in the selection of the measures in columns nine and ten, and the latest and best ones published up to the end of November 1885 have been given where possible. In the cases of binaries of quick revolution and stars with large proper motion, the position angles and distances have usually been roughly computed for the season in 1886 at which they appear in the lists where no very recent measures could be obtained of them by practised observers.

The eleventh column contains the magnitudes of the stars. In view of the great and, to beginners, most perplexing differences in the values assigned to the same star in the scales used by different astronomers, it has been thought well to adhere to one scale throughout the book, and for this purpose W. Struve's has been selected, as it is the one almost universally used by the great double star observers on the Continent and in America. The well-known English observer, Admiral Smyth, used a scale of his own, nearly agreeing with Sir John Herschel's, for the faint stars, and for the brighter he usually adopted the magnitudes assigned to the stars by Piazzi in his Palermo catalogue, estimating the magnitude of the

fainter component by a comparison of its light with that of the larger star, as given by Piazzi ; but where Piazzi gives both components of a double star Smyth usually adopted his magnitude of both. The following little table, which is copied from one in the *English Mechanic* for March 24, 1882, is the result of a direct collation of Smyth's magnitudes with Struve's, the magnitudes in Smyth above 9·5 being generally assumed from Piazzi, when the stars are found in the catalogue of that astronomer :—

Smyth Mag.		Struve Mag.	Smyth Mag.		Struve Mag.
9·5	=	8·78	13	=	10·65
10	=	9·18	14	=	10·36 <sup>1</sup>
11	=	10·10	15 }	=	10·99
12	=	10·17	16 }		

The following comparison by Sir John Herschel of his own magnitudes with those of Struve is copied from his table in vol. 38 of the 'Memoirs of the Royal Astronomical Society.' It must, however, be only considered to apply to his earliest catalogues of double stars, as he himself observes that the magnitudes in his later catalogues show a tendency to creep up in the scale, *i.e.* a 19th or 20th magnitude in his first catalogue would probably only be rated as a 16th in his sixth catalogue ; and it will be found that a star called by him a 20th magnitude in his later observations corresponds more nearly with 13·0 than 12·0 in Struve's<sup>2</sup> scale. The magnitudes down to the seventh are fairly accordant.

H. Mag.	Σ Mag.	H. Mag.	Σ Mag.	H. Mag.	Σ Mag.
8	= 7·30	11·5	= 9·60	15	= 10·87
8·5	= 7·70	12	= 9·80	16	= 11·13
9	= 8·10	12·5	= 10·00	17	= 11·38
9·5	= 8·50	13	= 10·18	18	= 11·61
10	= 8·80	13·5	= 10·36	19	= 11·82
10·5	= 9·10	14	= 10·54	20	= 12·00
11	= 9·30	14·5	= 10·71		

<sup>1</sup> See 'English Mechanic' *supra*.

<sup>2</sup> W. Struve's name is usually conventionally expressed by the Greek capital letter Σ, and his son's, O. Struve, the present director of the Imperial Russian Observatory at Pulkova, by OΣ.

In the present work the magnitudes of many of the brighter stars have been inserted from the Oxford photometrical determinations in vol. 47 of the 'Memoirs of the Royal Astronomical Society,'<sup>1</sup> and the others have been taken from the great work of the late Baron Dembowski, 'Misura Micrometriche di Stelle Doppie e Multiple,' which has lately been edited by Schiaparelli and O. Struve, in nearly every case where the stars are to be found in that work. In other cases they have been taken from the micrometrical observations of Mr. Burnham. The magnitudes of the red stars have been taken from Birmingham's 'Red Star Catalogue' and other sources. In the case of clusters the magnitudes of the brightest and faintest stars in each cluster have been set down.

In the last column are inserted the colours of the stars, which in the case of the double stars have been almost always taken from Dembowski. The discordances in the colours assigned by different astronomers to various stars, even where their observations have been made at the same epoch, are so very marked, that it has been thought advisable that these determinations should rest as far as possible on the authority of one observer only. This column contains also a few brief remarks on the character of some of the objects, which are supplemented in certain cases of interest by concise notes at the foot of each page. A considerable margin has been left for the observer's own notes. A list of about two hundred objects which can be seen with refractors of from four to seven inches aperture, has been added for the convenience of those amateurs who may possess telescopes of this size. A certain proportion only are visible, of course, with but four inches of aperture, but all should be seen with seven. Considerable care has been taken in the selection of the most interesting and beautiful objects within the grasp of telescopes of this size for the list, which is arranged in substantially the same way as the preceding one; the times of transit and the column containing the colours and remarks have, however, been omitted.

<sup>1</sup> The *Uranometria Nova Oxoniensis* was published too late to enable us to make use of the determinations of magnitude which it contains.

**Test Objects.**

The list of test-objects contains two hundred and fifty stars, which have been selected as tests of the quality of refracting telescopes of from two to seven inches aperture. The stars have been divided into three kinds of tests, viz. dividing tests, defining tests, and space-penetrating tests. Twelve double stars have been arranged, in descending order of difficulty, as tests of the separating power of telescopes of the aperture given. The first star of these twelve should be easily separated with the aperture employed, as high a magnifying power as the telescope and the air will bear being used, while the last three or four stars will be found to be just at the limit of the separating power of the aperture, this being expressed by the fraction  $\frac{4.56''}{\alpha}$ .<sup>1</sup> These are followed by a list of six pairs arranged in the same order, which are intended to serve as tests of the defining power of the telescope, its freedom from spherical aberration, and of the perfection of its figure; each pair consisting of a bright star and a faint and comparatively close companion. The last twelve stars (arranged, as the others, approximately in order of difficulty) are designed to serve as tests of the light-grasping power of the instrument and the acuteness of the observer's eye.

The list of the radiant points of some of the most marked showers of shooting stars gives the date of each shower, the Right Ascension and Declination of the radiant point, and the Greenwich mean time of its rising and southing.

**Lunar Craters.**

The lists of the approximate longitudes and latitudes of over one hundred lunar craters, arranged alphabetically and in order of longitude, beginning at the western limb, is intended to facilitate the employment of the table giving the position of the Lunar

<sup>1</sup> Dawes, *Memoirs of the Royal Astronomical Society*, vol. xxxv. p. 158. Dallmeyer preferred  $\frac{4.33''}{\alpha}$ .  $\alpha$  = aperture of telescope in inches.



Terminator at midnight on each day of the year. The number in ( ) immediately following the name of each formation refers to the corresponding number on the map of the moon, in the fourth edition of 'Celestial Objects for Common Telescopes,' the numbers enclosed in [ ] denoting formations near those bearing the number assigned to them by Mr. Webb. The figures i, ii, iii, iv refer to the quadrant in which the crater is situated.

The table giving the position of the Lunar Terminator at midnight for every night in the year gives the selenographical longitude of the point where the terminator (or boundary between light and darkness) crosses the moon's equator (which it does nearly at right angles) at that time. As the terminator moves towards the east at the rate of  $30^{\circ}5'$  (approximately) per hour,<sup>1</sup> we can easily find its position at any other time; *e.g.* by the table we find the longitude of the terminator at midnight on March 8, 1886, to be  $49^{\circ}26'$  west longitude, and we wish to ascertain what will its position be at 8 A.M. the next morning (March 9)? Subtracting  $4^{\circ}4'$  ( $30^{\circ}5' \times 8$ ), as the terminator moves towards the east, we find the longitude of the terminator at 8 in the morning on March 9 is  $45^{\circ}22'$  west. In the same way we wish to know when the crater Linné will be on the terminator in January 1886. Turning to the alphabetical table of the longitudes and latitudes of the craters, we find the longitude of Linné to be  $11^{\circ}33'$  west, and we find by the tables giving the position of the Lunar Terminator that the moon's terminator was in  $10^{\circ}44'$  west at midnight on January 11; the difference is  $0^{\circ}49'$  west, or the time of sunrise on Linné will be about  $10^{\text{h}}22^{\text{m}}$  P.M. (that is, 12 hours less  $49 \times 2$  min.) Similarly, we find that the longitude of the evening terminator at midnight on January 26 was  $8^{\circ}9'$ , differing  $3^{\circ}24'$  from that of Linné, so that sunset on Linné will take place about  $5^{\text{h}}20^{\text{m}}$  P.M. on that evening. Conversely, we observe that the longitude of the evening terminator at midnight on February 11, 1886, is given in the tables at  $6^{\circ}15'$  east, and on referring to the list of craters arranged in order of longitude, we find Mösting, Moretus, and Archimedes A to be near the terminator at that time, being numbers 211, 262, and a

<sup>1</sup> Roughly speaking,  $1^{\circ}$  for two hours, or  $1'$  for two minutes.

crater very near 120 on Webb's map. On referring to that map, and drawing an imaginary line through these craters, we are enabled to see what other formations are on the terminator about that hour.

### **Variable Stars, &c.**

A list of the maxima and minima of twelve remarkable variable stars, with their places, and a table showing the observable heliocentric maxima of Algol and of the somewhat similar variable star  $\delta$  Libræ for 1886, will, it is hoped, be found of use to those amateurs who are interested in the study of such phenomena.

A short account of the periodical comets expected in 1886 and of those which, though discovered in 1885, pass their perihelia in 1886, closes the work.

### **Precession of the Equinoxes.**

Owing to precession, the apparent places of the stars alter continually from year to year, both in Right Ascension and in Declination; the change is fortunately very small, and may generally be neglected, as even after several years it is seldom sufficient to prevent a star being found in the field of the telescope. Its amount varies in different parts of the heavens, and the formulæ for its computation are somewhat complex, but the tables given below will enable the observer to correct the places of the stars for this quantity without any very serious error, except in the case of circum-polar stars.

The precession of the Equinoxes is caused by a slow gyratory or twisting motion of the pole of the earth round the pole of the Ecliptic, resembling the movement of a spinning top just before it is going to fall. One twist of the axis occupies about 25,700 years, and during this time it carries the position of the vernal Equinox or first point of Aries backwards completely round the Ecliptic, in a direction contrary to the apparent motion of the sun, thereby continually increasing the apparent Right Ascension of all the heavenly bodies, and also affecting their Declinations; the effect

on our reckoning of time is the same as if we had gained one day in 25,700 years, or  $\frac{1}{25700}$  of a day per annum. This amounts to about 3.1 seconds, which is about the annual amount of precession for stars near the Equator.

The effect of precession in Right Ascension is always additive (except in the case of a few stars very near the poles of the Ecliptic). Its amount varies with the Declination of the stars and with their Right Ascension, and is roughly given in the following table :—

*Annual Value of Precession in Right Ascension.*

Declination	From 0 <sup>h</sup> to 6 <sup>h</sup> Right Ascension	From 6 <sup>h</sup> to 12 <sup>h</sup> Right Ascension	From 12 <sup>h</sup> to 18 <sup>h</sup> Right Ascension	From 18 <sup>h</sup> to 24 <sup>h</sup> Right Ascension
20° South Equator	3 <sup>s</sup> .1 decreasing to 2 <sup>s</sup> .6 3.1 secs.	2 <sup>s</sup> .6 increasing to 3 <sup>s</sup> .1 3.1 secs.	3 <sup>s</sup> .1 increasing to 3 <sup>s</sup> .6 3.1 secs.	3 <sup>s</sup> .6 decreasing to 3 <sup>s</sup> .1 3.1 secs.
20° North	3 <sup>s</sup> .1 increasing to 3 <sup>s</sup> .6	3 <sup>s</sup> .6 decreasing to 3 <sup>s</sup> .1	3 <sup>s</sup> .1 decreasing to 2 <sup>s</sup> .6	2 <sup>s</sup> .6 increasing to 3 <sup>s</sup> .1
40° North	3 <sup>s</sup> .1 increasing to 4 <sup>s</sup> .2	4 <sup>s</sup> .2 decreasing to 3 <sup>s</sup> .1	3 <sup>s</sup> .1 decreasing to 1 <sup>s</sup> .9	1 <sup>s</sup> .9 increasing to 3 <sup>s</sup> .1
50° North	3 <sup>s</sup> .1 increasing to 4 <sup>s</sup> .7	4 <sup>s</sup> .7 decreasing to 3 <sup>s</sup> .1	3 <sup>s</sup> .1 decreasing to 1 <sup>s</sup> .5	1 <sup>s</sup> .5 increasing to 3 <sup>s</sup> .1
60° North	3 <sup>s</sup> .1 increasing to 5 <sup>s</sup> .4	5 <sup>s</sup> .4 decreasing to 3 <sup>s</sup> .1	3 <sup>s</sup> .1 decreasing to 0 <sup>s</sup> .8	0 <sup>s</sup> .8 increasing to 3 <sup>s</sup> .1

The precession in Declination is additive from 18 hours Right Ascension to 6 hours, and subtractive from 6 hours to 18 hours. It attains its maximum, about 20 seconds of arc per annum, with objects situated at 12 hours Right Ascension and 24 hours Right Ascension, and falls to a minimum of zero at 6 hours and 18 hours. The following table will give an idea of the change produced by it, but

*Annual Value of Precession in Declination.*

Right Ascension		Annual Precession	Right Ascension		Annual Precession
h. m.	and h. m.	"	h. m.	and h. m.	"
0 0	and 24 0	+ 20.0	12 0	and 12 0	- 20.0
1 0	23 0	19.4	13 0	11 0	19.4
2 0	22 0	17.4	14 0	10 0	17.4
2 30	22 30	15.9	14 30	9 30	15.9
3 0	21 0	14.2	15 0	9 0	14.2
3 30	20 30	12.2	15 30	8 30	12.2
4 0	20 0	10.0	16 0	8 0	10.0
4 30	19 30	7.7	16 30	7 30	7.7
5 0	19 0	5.2	17 0	7 0	5.2
5 30	18 30	2.6	17 30	6 30	2.6
6 0	18 0	0.0	18 0	6 0	0.0

it must be remembered that the table refers to stars with North Declination, and that an addition to the Declination of a northern star corresponds to a subtraction from that of a southern star, and

*vice versa*. It will be observed that at the utmost it can only change the declination of any object by 20'' per annum, so that by the year 1892 the change will only have amounted to 2', or about three times the apparent diameter of Jupiter when at opposition.

### Future Use of the Tables.

The times of transit of all the objects in this book are given for the year 1886, but by a very simple addition they are equally available for all succeeding years. We have only to add one minute for every year, except at leap-years, when we make a deduction. For example :—

1887. Add 1 minute to the times given.

1888. Add 2 minutes before February 28 ; after that date deduct 2 minutes.

1889. Deduct 1 minute.

1890. Add 0 minute.

1891. Add 1 minute.

It will be perceived that, although for convenience we abruptly add 1 minute on January 1, the gain is really a continuous one throughout the four years preceding the leap-year. At leap-year we suddenly lose this 4 minutes (or rather 3<sup>m</sup> 56<sup>s</sup>) owing to the introduction of February 29.

---

It was originally intended that this work should be published in an annual form under the name of 'Clark's Star Guide,' as a companion volume to Mr. Latimer Clark's annual Transit Tables. As the work progressed it was found difficult to condense the matter satisfactorily into so small a page, and it was therefore thought preferable to issue it in the present more permanent form. The tables giving the daily position of the Lunar Terminator and other matters will be continued in future editions of Clark's Transit Tables.



A LIST OF  
THE MOST REMARKABLE CELESTIAL OBJECTS  
VISIBLE WITH SMALL TELESCOPES.

NAME OF OBJECT	Visible	R.A.		Decl.	1		11		21		31		Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h.	m.		h.	m.	h.	m.	h.	m.	h.	m.				
Sidereal time at 9 P.M.	—	—	—	—	3	46	4	25	5	5	4	44	—	—	—	—
ψ Aquarii . . . . .	Nov.—Jan.	23	9'9	— 9 43	4	25	3	46	5	5	4	44	49'4	312	4.0, 8.5	Orange, sky blue (1).
ο Cephei . . . . .	July—Mar.	23	13'9	+67 29	4	29	3	50	3	10	2	31	2'6	193	5.2, 7.6	Deep yellow, turquoise. Binary.
Cluster in Cepheus . . . . .	Aug.—Feb.	23	19'2	+60 58	4	34	3	55	3	16	2	36	—	—	9-11	6' diam. Triangular-shaped (2).
77 Pegasi . . . . .	Dec.—Feb.	23	37'5	+ 9 42	4	52	4	13	3	34	2	54	—	—	5.0	Red.
107 Aquarii . . . . .	Dec.—Jan.	23	40'1	+19 19	4	55	4	16	3	36	2	57	5'6	140	5.3, 6.5	White, bluish. Colours vary.
σ Cassiopeiæ . . . . .	Aug.—Feb.	23	53'2	+55 6	5	8	4	29	3	49	3	10	2'9	327	4.8, 7.1	Greenish white, blue.
Pair in Cassiopeiæ . . . . .	Aug.—Feb.	23	55'5	+59 43	5	10	4	31	3	52	3	12	58'9	270	7.0, 7.7	Ruby, deep blue. 7.0 variable.
Pair in Cassiopeiæ* . . . . .	Aug.—Feb.	0	0'3	+57 48	5	15	4	36	3	57	3	17	1'7	311	6.0, 7.0	Yellow, olive. Binary (3).
35 Piscium . . . . .	Dec.—Feb.	0	9'1	+ 8 11	5	24	4	45	4	5	3	26	11'5	149	5.8, 7.2	White, deep blue (4).
α Cassiopeiæ . . . . .	Aug.—Feb.	0	34'0	+55 55	5	49	5	10	4	30	3	51	62'4	280	2.4, 9'5	Pale rose, blue. 2.4 variable?
Great Neb. in Androm. . . . .	Aug.—Feb.	0	36'3	+40 35	5	51	5	12	4	32	3	53	—	—	—	Hazy 5½ mag. star to naked eye
γ Cassiopeiæ . . . . .	Aug.—Feb.	0	42'1	+57 13	5	57	5	18	4	38	3	59	4'9	175	3'5, 7.3	Yellow, rose. Binary (6). (15)
36 Andromedæ* . . . . .	Sept.—Feb.	0	48'9	+23 1	6	4	5	24	4	45	4	6	1'5	2	6.0, 6.4	Both golden. Binary (7).
77 Piscium . . . . .	Dec.—Feb.	0	59'9	+ 4 18	6	15	5	35	4	56	4	17	32'8	83	6'1, 6'8	White, pale lilac.
ζ Piscium . . . . .	Dec.—Feb.	1	7'7	+ 6 57	6	23	5	43	5	4	4	25	23'8	64	4.2, 5'8	White, pale rose. 4.2 variable.
35 Cassiopeiæ . . . . .	Aug.—Feb.	1	13'5	+64 4	6	28	5	49	5	10	4	30	52'5	350	6'1, 8'0	White, deep red.
Nebula in Triangulum . . . . .	Oct.—Feb.	1	27'4	+30 4	6	42	6	3	5	23	4	44	—	—	—	40' by 12'. Faint (8).
Cluster in Cassiopeiæ . . . . .	Aug.—Feb.	1	38'2	+60 40	6	53	6	14	5	34	4	55	—	—	8-11	Fine field. Several pairs (9).
γ Arietis . . . . .	Nov.—Feb.	1	47'2	+18 44	7	2	6	22	5	43	5	4	8'8	179	4.0, 4'2	Both white (10).
α Piscium . . . . .	Nov.—Feb.	1	56'1	+ 2 13	7	11	6	31	5	52	5	13	3'0	323	4'1, 5'4	Green, ash green.
γ Andromedæ . . . . .	Aug.—Feb.	1	56'9	+41 46	7	11	6	32	5	53	5	14	10'4	63	2'1, 6'1	Golden, ultramarine (11).
ι Trianguli . . . . .	Nov.—Feb.	2	5'7	+29 46	7	20	6	41	6	2	5	22	3'6	78	5'1, 6'7	Golden, blue.
Clusters in Perseus . . . . .	Aug.—Feb.	2	11'2	+56 38	7	26	6	46	6	7	5	28	—	—	6'5-13	Visible to naked eye (12).
ι Cassiopeiæ* . . . . .	July—Mar.	2	19'7	+66 53	7	34	6	55	6	16	5	36	7'7	108	5'0, 8'1	Yellowish white, deep blue (13).

(1) Moving together through space.  
 (2) The brightest star is red.  
 (3) Period about 105 years; pair in field, 7.0, 8.5; 299°: 80' 5".  
 (4) 38 Piscium follows nearly on the parallel at 2m 25'; 6.5, 7.2; 238°: 4" 3", white, ashy.  
 (5) The 'Queen of the Nebulae,' oval, very extended in great telescopes, milky white. A round white nebula, 30" diameter, 20" south of the great one, and a faint oval one north preceding.  
 (6) Period probably 140 years; mass of system eight times greater than that of the Sun; distance, 1,340,000 times that of the Earth from the Sun.  
 (7) Period 316 years.  
 (8) Requires low powers; a 6½ mag. star in the field, north.  
 (9) Another cluster, triangular-shaped, 2m preceding and 46' north of this.  
 (10) A 9 mag. star at 85°: 229'.  
 (11) 6'1 is an excessively close pair, distance 0" 36'.  
 (12) Two magnificent masses, forming the 'sword-handle' of Perseus, from the 5.0 mag., ruddy violet.

NAME OF OBJECT	Visible	R.A.		Decl.	1		11		21		31		Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h. m.	o		h. m.	h. m.	h. m.	h. m.	h. m.	h. m.						
Sidereal time at 9 P.M.																
30 Arietis . . . . .	Nov.—Feb.	2 30.3	+ 24 9	—	3 46	5 5	5 44	—	—	—	—	—	—	—	—	White, deep blue.
γ Ceti . . . . .	Nov.—Jan.	2 37.4	+ 2 45	—	7 45	7 5	6 26	5 47	5 47	5 47	5 47	38.6	273	5.4, 6.6	—	Whitish yellow, ash.
α Ceti . . . . .	Nov.—Jan.	2 56.3	+ 3 38	—	7 52	7 12	6 33	5 54	5 54	5 54	5 54	2.8	290	3.4, 7.1	—	Orange (1).
Cluster in Perseus* . . . . .	Aug.—Feb.	3 6.9	+ 46 47	—	8 21	7 42	6 52	6 13	6 13	6 13	6 13	—	—	10-12	—	Large stars arranged in lines.
P. iii. 98 Eridani . . . . .	Dec.—Feb.	3 30.9	+ 0 13	—	8 45	8 6	7 27	6 47	6 47	6 47	6 47	6.2	246	6.3, 8.4	—	Deep yellow, blue.
Nebula in Eridanus* . . . . .	Dec.—Feb.	3 35.1	- 18 58	—	8 49	8 10	7 31	6 51	6 51	6 51	6 51	—	—	—	—	Pale, but distinct. 60" diam.
γ Tauri . . . . .	Nov.—Feb.	3 40.7	+ 23 45	—	8 55	8 16	7 36	6 57	6 57	6 57	6 57	147.3	289	3.0, 6.0	—	White, pearly (2).
32 Eridani . . . . .	Dec.—Feb.	3 48.6	- 3 18	—	9 3	8 24	7 44	7 5	7 5	7 5	7 5	6.8	347	5.0, 6.7	—	Deep yellow, violet.
ε Persei* . . . . .	Oct.—Feb.	3 50.2	+ 39 41	—	9 4	8 25	7 46	7 7	7 7	7 7	7 7	8.7	10	3.0, 7.9	—	Greenish yellow, ashy blue.
Cluster in Perseus . . . . .	Aug.—Feb.	4 6.6	+ 50 57	—	21	8 42	8 2	7 23	7 23	7 23	7 23	—	—	9-11	—	Large stars in curves.
δ Tauri . . . . .	Oct.—Feb.	4 13.3	+ 27 5	—	9 28	8 48	8 9	7 30	7 30	7 30	7 30	53.6	246	5.0, 8.0	—	Light red, sky blue.
ι Camelopardi . . . . .	Aug.—Feb.	4 23.0	+ 53 40	—	9 37	8 58	8 19	7 39	7 39	7 39	7 39	10.2	397	5.3, 6.3	—	Deep greenish yellow, green.
α Tauri* (Aldebaran) . . . . .	Oct.—Feb.	4 29.4	+ 16 17	—	9 44	9 4	8 25	7 46	7 46	7 46	7 46	115.1	34	1.1, 10.3	—	Rose, blue (3).
ω Aurigæ . . . . .	Oct.—Mar.	4 51.5	+ 37 43	—	10 6	9 26	8 47	8 8	8 8	8 8	8 8	6.0	353	5.1, 7.7	—	White, reddish blue.
Red star in Auriga . . . . .	Oct.—Mar.	4 52.6	+ 39 29	—	10 7	9 27	8 48	8 9	8 9	8 9	8 9	—	—	6.4	—	Red.
β Camelopardi . . . . .	July—Mar.	4 53.3	+ 60 16	—	10 7	9 28	8 49	8 9	8 9	8 9	8 9	80.3	208	4.0, 7.2	—	Deep yellow, white.
14 Aurigæ . . . . .	Oct.—Mar.	5 8.0	+ 32 33	—	10 23	9 43	9 3	8 24	8 24	8 24	8 24	14.7	225	5.1, 7.5	—	Deep yellow, olive.
β Orionis (Rigel) . . . . .	Dec.—Feb.	5 9.1	- 8 20	—	10 22	9 44	9 4	8 25	8 25	8 25	8 25	9.5	201	1.0, 7.8	—	Yellowish white, blue.
III Tauri . . . . .	Oct.—Feb.	5 17.7	+ 17 17	—	10 32	9 52	9 13	8 34	8 34	8 34	8 34	74.8	271	5.7, 8.3	—	Yellow, lilac (4).
δ Orionis . . . . .	Dec.—Mar.	5 26.2	- 0 23	—	10 40	10 1	9 22	8 42	8 42	8 42	8 42	52.5	359	2.0, 6.6	—	Blue, pearly.
θ Orion. and Nebula . . . . .	Dec.—Mar.	5 29.7	- 5 28	—	10 44	10 4	9 25	8 46	8 46	8 46	8 46	8.8	33	7.1, 7.9	—	Both blue, A-B. See (5).
ζ Orionis* . . . . .	Dec.—Mar.	5 35.0	- 2 0	—	10 49	10 10	9 30	8 51	8 51	8 51	8 51	2.6	152	2.3, 5.8	—	White, blue (6).
γ Leporis . . . . .	Jan.—Feb.	5 39.7	- 22 29	—	10 54	10 14	9 35	8 56	8 56	8 56	8 56	93.7	350	4.0, 6.5	—	Yellow, garnet (7).
Cluster in Auriga . . . . .	Oct.—Mar.	5 44.8	+ 32 31	—	10 59	10 19	9 40	9 1	9 1	9 1	9 1	—	—	9-11	—	500 stars. Magnificent.

(1) A blue 6.5 mag., 16' north, with faint pair bearing.  
 (2) A blue 7.6 mag. at 312° : 180"/9, and a blue 8.2 mag. at 295° : 190"/8, the three forming a neat triangle.  
 (3) Distance of Aldebaran from the solar system 400,000 times greater than that of the Earth from the Sun.  
 (4) Rectilinear motion.  
 (5) The Trapezium in the midst of the great Nebula, one of the most marvellous objects in the heavens. The Trapezium consists of four bright stars and two very faint ones : the brightest star of the Trapezium, C, 4.9 mag., bright yellow, is 13 1/2" distant from A (the 7.1 mag. given above), on an angle of 31° ; the fourth bright star, D, 6.8 mag., blue, is 13 1/4" from C, on an angle of 61°. The nebula itself is of a greenish hue, and in large telescopes extends over several degrees. It contains several variable stars, and probably consists of various glowing gases.  
 (6) A 9.5 mag. at 9° : 57"/2.  
 (7) A small star near the 6.5 mag. Common proper motion.

FEBRUARY

FEBRUARY (Mean Time of Transit at Greenwich).

NAME OF OBJECT	Visible	R.A.	Decl.	1	11	21	28	Dis- tances	Posi- Angles	Mag.	COLOURS AND REMARKS
		h. m.	o /	h. m.	h. m.	h. m.	h. m.	"	o	—	—
Sidereal time at 9 P.M.	—	0 36.3	+40 35	3 49	3 10	2 31	7 34	—	—	—	—
Great Neb. in Andromeda	Aug.—Feb.	0 42.1	+57 13	3 55	3 16	2 36	2 9	4.9	175	3.5, 7.3	45' by 15' in a 3 inch (1). Yellow, rose (1). Copper red.
η Cassiopeiæ	Aug.—Mar.	1 5.9	+44 42	4 19	3 39	3 0	2 33	—	—	6.5	Brilliant group, surrounding φ.
B.A.C. 352 Andromedæ	Aug.—Mar.	1 12.0	+58 11	4 25	3 45	3 6	2 39	—	—	7-9.5	Fan-shaped. Contains a red star
Cluster in Cassiopeia	Aug.—Mar.	1 25.9	+60 4	4 39	3 59	3 20	2 53	—	—	7.5-10	White, yellow. [(2).
χ <sup>1</sup> Ceti	Nov.—Feb.	1 36.1	-11 54	4 49	4 10	3 30	3 3	3.5	88	5.9, 7.0	Orange, turquoise.
Pair in Cassiopeia	Aug.—Mar.	1 43.0	+64 17	4 56	4 16	3 37	3 10	34.9	34	6.0, 8.2	Golden, deep green.
P. i. 179 Arietis	Nov.—Mar.	1 43.7	+21 42	4 56	4 17	3 38	3 10	2.7	168	6.1, 7.0	Deep yellow, yellow.
56 Andromedæ	Aug.—Feb.	1 49.3	+36 41	5 2	4 23	3 43	3 16	183.7	121	5.7, 5.9	Golden, ultramarine (1).
γ Andromedæ	Aug.—Feb.	1 56.9	+41 50	5 10	4 30	3 51	3 23	10.4	63	2.1, 6.1	White, pale rose. Pair α, φ.
59 Andromedæ	Sep.—Mar.	2 4.1	+38 30	5 17	4 37	3 58	3 31	16.5	36	6.0, 6.7	Yellow, sapphire (3).
66 Ceti	Oct.—Mar.	2 6.9	-2 56	5 20	4 40	4 1	3 33	15.6	229	5.9, 7.5	Contain several red stars (1).
Clusters in Perseus	Aug.—Mar.	2 11.2	+56 38	5 24	4 45	4 5	3 38	—	—	6.5-13	Reddish, blue. Max. Jan. 5 (4).
o Ceti (Mitra)	Oct.—Mar.	2 13.6	-3 30	5 26	4 47	4 8	3 40	115.6	83	Var. 9.5	Visible to naked eye. 15' diam.
Cluster in Perseus	Sep.—Mar.	2 34.7	+42 15	5 47	5 8	4 29	4 1	—	—	8-10	Round, faint; 9 m. star at 130° :2.
Nebula in Cetus *	Oct.—Mar.	2 36.8	-0 28	5 49	5 10	4 31	4 3	—	—	—	Golden, blue (5).
Perseid *	Sep.—Mar.	2 42.4	+55 25	5 55	5 16	4 36	4 9	28.4	300	4.0, 8.0	White.
Pair in Cetus	Oct.—Mar.	3 3.1	+6 59	6 16	5 36	4 57	4 29	81.2	163	7.0, 7.2	White, ash.
Pair in Perseus	Oct.—Mar.	3 9.8	+40 3	6 22	5 43	5 4	4 36	3.4	28	6.4, 7.2	White, ashy blue.
Pair in Camelopardus *	Sep.—Mar.	3 20.9	+58 58	6 33	5 54	5 15	4 47	2.6	64	6.4, 7.6	Orange red, blue.
Pair in Camelopardus	Sep.—Apr.	3 33.3	+59 36	6 46	6 10	5 27	5 0	55.6	34	6.0, 9.0	Red, deep blue (6).
Pair in Perseus	Oct.—Mar.	3 36.5	+38 1	6 49	6 13	5 31	5 3	30.2	87	6.7, 7.5	Fine orange.
Red star in Camelopardus	Sep.—Apr.	3 38.9	+65 10	6 51	6 12	5 33	5 5	—	—	6.0	White, blue (7).
P. iii. 213 Tauri	Nov.—Mar.	3 54.1	+22 53	7 6	6 27	5 48	5 20	7.3	128	6.5, 7.7	

(1) See January list. A new star has recently appeared in the Great Nebula in Andromedæ, near the nucleus, but is now rapidly fading.  
 (2) A 9 mag. star in α, γ quadrant, from which the cluster takes its divergence.  
 (3) Moving together through space; common proper motion 0".5 annually.  
 (4) See list of Variable Stars.  
 (5) Forms with three other stars a miniature of Jupiter and his four satellites.  
 (6) Recilinear motion.  
 (7) A 9.0 mag. var. <sup>9</sup>, at 241° : 58" 1.



FEBRUARY

NAME OF OBJECT	Visible	R.A.		Decl.		1		11		21		28		Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h.	m.	o.	'	h.	m.	h.	m.	h.	m.	h.	m.				
Sidereal time at 9 P.M.																	
Cluster in <b>Persæus</b>	Oct.—Mar.	4	1'5		+49 12	4	48	6	27	7	7	7	34	—	—	—	5' diam. Ring of stars round it.
Pair in <b>Taurus</b>	Nov.—Mar.	4	9'4		+5 34	7	22	6	35	5	55	5	36	—	—	—	Whitish yellow, rosy white.
x <b>Tauri</b>	Nov.—Mar.	4	15'6		+25 21	7	28	6	49	6	9	5	42	65'5	315	6'2, 6'7	White, reddish blue.
γ <b>Tauri</b>	Nov.—Mar.	4	35'4		+22 44	7	48	7	8	6	29	6	2	19'2	25	57, 8'2	White, bluish white.
55 <b>Eridani</b>	Dec.—Mar.	4	38'1		— 9 0	7	50	7	11	6	32	6	4	62'9	212	5'0, 7'3	Rosy white, white.
P. iv. 255-7 <b>Tauri</b>	Nov.—Mar.	4	52'5		+14 23	8	5	7	25	6	46	6	19	9'0	316	6'0, 6'3	Bright green, bright blue (1).
11-12 <b>Camelopardi</b>	Oct.—Apr.	4	56'3		+58 52	8	8	7	29	6	50	6	22	39'2	305	5'3, 6'7	White, orange.
Pair in <b>Auriga</b> *	Nov.—Mar.	5	2'6		+37 9	8	15	7	35	6	56	6	28	180'8	8	4'8, 5'9	Deep red, deep green.
ι <b>Leporis</b> *	Dec.—Mar.	5	7'0		—12 0	8	19	7	40	7	1	6	33	1'6	221	6'8, 7'1	White, pale violet (2).
β <b>Orionis</b> ( <i>Rigel</i> ).	Dec.—Mar.	5	9'1		— 8 20	8	21	7	42	7	3	6	35	12'6	336	4'5, 10'0	Yellowish white, blue (3).
Cluster in <b>Auriga</b>	Nov.—Mar.	5	21'8		+35 48	8	34	7	55	7	15	6	48	9'5	201	1'0, 7'8	Arranged as an oblique cross.
δ <b>Orionis</b>	Dec.—Mar.	5	26'2		— 0 23	8	38	7	59	7	20	6	52	—	—	7'5-11	Blue, pearly (3).
λ <b>Orionis</b>	Dec.—Mar.	5	28'8		+ 9 52	8	41	8	2	7	22	6	55	52'5	359	2'0, 6'6	White, deep blue.
θ <sup>o</sup> <b>Orionis</b> and <b>Nebula</b>	Dec.—Mar.	5	29'7		— 5 28	8	42	8	2	7	23	6	50	4'3	45	4'2, 6'2	Deep yellow, whitish blue (4).
σ <b>Orionis</b> *	Dec.—Mar.	5	33'0		— 2 40	8	45	8	6	7	26	6	59	52'6	92	3'5, 5'5	White blue (5).
P. v. 214 <b>Tauri</b>	Nov.—Mar.	5	40'9		+24 39	8	53	8	14	7	34	7	7	11'1	235	3'9, 9'5	Deep yellow, red.
Cluster in <b>Auriga</b>	Nov.—Mar.	5	44'8		+32 31	8	57	8	18	7	38	7	11	94'2	166	6'5, 7'0	24' diam. including outliers.
Cluster in <b>Gemini</b>	Nov.—Mar.	6	1'8		+24 22	9	14	8	35	7	55	7	28	—	—	8-11	Visible to naked eye.
Pair in <b>Orion</b>	Dec.—Mar.	6	12'9		+13 29	9	25	8	46	8	6	7	39	73'3	44	6'5, 7'0	White, red.
ι1 <b>Monocerotis</b>	Dec.—Apr.	6	23'3		— 6 57	9	35	8	56	8	17	7	49	7'4	132	5'7, 5'7	White, whitish yellow (6).
Cluster in <b>Monoceros</b>	Dec.—Apr.	6	26'2		+ 4 57	9	38	8	59	8	20	7	52	—	—	6-11	Scattered; surrounds 12 Mon.
56 <b>Aurigæ</b>	Nov.—Apr.	6	38'5		+43 41	9	50	9	11	8	32	8	4	47'8	22	5'5, 8'0	Yellow, turquoise (7).
Cluster in <b>Canis Major</b>	Dec.—Mar.	6	42'1		—20 38	9	54	9	15	8	35	8	8	—	—	8-10	Visible to naked eye.
ζ <b>Geminorum</b>	Dec.—Apr.	6	57'4		+20 44	10	9	9	30	8	51	8	23	93'6	352	4'0, 7'2	Deep yellow, deep blue (8).

(1) Also called 26 (Bode) **Orionis**. A 9<sup>o</sup> mag., purple, at 88° 54''/7'.  
 (2) A very red 6 mag. (variable) 57 seconds preceding on an angle of 274°.  
 (3) See Jan. list; 7'8 mag. companion of **Rigel** is an excessively close double, 0''/2.  
 (4) A-C 08°; 128''/8; 7'6 mag. These are the three stars in a line, above the fish's mouth in the Great Nebula. The 3'5 mag. is 135''/2 distant from the brightest star of the Trapezium in the direction of 134°.  
 (5) A 6'8 mag., blue, at 85° 12''/8 from A, and a 6'3 mag., reddish, at 61° 44''/6 from 6.  
 (6) B double, 6'1 mag., white, 103° 21''/5. A 7 mag. 250''/4, 4.  
 (7) Moving, owing to proper motion of A.  
 (8) A faint star, 10 5 mag., at 84° 0' 87''/2.

MARCH (Mean Time of Transit at Greenwich).

MARCH

NAME OF OBJECT	Visible	R.A.	Decl.	1	11	21	31	Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
Sidereal time at 9 P.M.	—	h. m.	o	h. m.	h. m.	h. m.	h. m.	"	o	—	—
Cluster in Perseus . . .	Oct.—Mar.	4 12'3	—	7 38	8 18	8 57	9 37	—	—	6'5-10	Radiated. 10' diam. (1).
Pair in Perseus . . .	Nov.—Mar.	4 16'9	+34 3	5 39	5 0	4 20	3 37	—	—	6'3, 7'5	White. Fine field.
62 Tauri . . .	Nov.—Mar.	4 17'1	+24 2	5 39	5 0	4 21	3 41	28'9	290	6'0, 8'0	White, blue. Fine field.
4' Tauri . . .	Nov.—Mar.	4 18'6	+22 0	5 41	5 2	4 22	3 43	340'4	173	4'6, 5'7	White, ashy yellow.
88 Tauri . . .	Nov.—Mar.	4 29'4	+9 56	5 52	5 12	4 33	3 54	69'2	299	4'0, 7'5	White, reddish yellow.
Pair in Orion . . .	Dec.—Mar.	4 54'6	+3 27	6 17	5 37	4 58	4 19	21'0	260	6'0, 7'0	White, deep green.
P. iv. 278 Orionis . . .	Dec.—Mar.	4 56'1	+1 27	6 18	5 39	5 0	4 20	14'1	49	6'3, 7'7	White, yellow.
Pair in Camelopardus . . .	Oct.—Apr.	5 1'7	+69 42	6 24	5 44	5 5	4 26	5'3	221	7'0, 8'2	Yellow, blue.
ρ' Orionis * . . .	Dec.—Mar.	5 7'3	+2 44	6 29	5 50	5 11	4 31	6'8	65	4'8, 8'5	Deep yellow, blue (2).
23 Orionis . . .	Dec.—Mar.	5 16'8	+3 26	6 39	6 0	5 20	4 41	32'1	28	5'1, 6'6	White, deep blue.
118 Tauri . . .	Nov.—Mar.	5 22'2	+25 3	6 44	6 5	5 26	4 46	4'7	199	5'5, 6'4	White, green (3).
Nebula in Taurus . . .	Nov.—Mar.	5 27'6	+21 56	6 50	6 10	5 31	4 52	—	—	—	Faint, 5½' by 3½' (4).
Cluster in Auriga . . .	Nov.—Mar.	5 28'7	+34 4	6 51	6 11	5 32	4 53	—	—	8'0-11	20' diam.; star-shaped.
γ Orionis . . .	Dec.—Mar.	5 29'8	-6 0	6 52	6 13	5 33	4 54	11'4	142	3'7, 7'7	Yellow, blue (5).
Nebulae in Orion * . . .	Dec.—Mar.	5 40'9	0 0	7 3	6 24	5 44	5 5	—	—	—	Double nebula, 7' diam. (6).
α Orionis (Betelgeux) . . .	Dec.—Mar.	5 49 0	+7 23	7 11	6 32	5 52	5 13	174'7	152	1'0, 8'9	Topaz, blue (7).
41 Aurigæ . . .	Nov.—Apr.	6 3'0	+48 44	7 25	6 46	6 5	5 27	7'9	354	5'8, 6'7	White, ashy (8).
8 Monocerotis . . .	Dec.—Mar.	6 17'7	+4 39	7 40	7 0	6 21	5 42	13'7	27	4'2, 6'2	Golden, lilac (9).
20 Monocerotis . . .	Dec.—Apr.	6 25'6	+17 52	7 48	7 8	6 29	5 50	19'9	210	6'1, 7'0	White, greenish white.
Red star in Auriga . . .	Dec.—Apr.	6 28'7	+38 32	7 51	7 11	6 32	5 53	—	—	6'0	Fine red (10).
γ Canis Majoris . . .	Dec.—Mar.	6 31'4	-18 34	7 53	7 14	6 35	5 55	17'5	261	6'0, 7'5	Pale red, grey.
38 Geminorum . . .	Dec.—Mar.	6 48'2	+13 19	8 10	7 31	6 51	6 12	6'3	162	5'4, 7'8	White, deep rose (11).
P. vi. 301 Lynceis . . .	Dec.—Apr.	6 56'6	+52 56	8 18	7 39	7 0	6 20	3'2	156	6'5, 6'7	White.
Cluster in Monoceros . . .	Dec.—Mar.	6 57'3	-8 11	8 19	7 40	7 0	6 21	—	—	8'0, 12'0	Brilliant mass (12).

(1) Like a badge of knighthood. Contains pair 6'5, 7'0; 327°; 74'6; red, white.  
 (2) A coarse pair in field, s. s. s.  
 (3) Common proper motion.  
 (4) Messier's, the so-called 'Crab' nebula, non-resolvable.  
 (5) An 8½ mag. at 100"8; 49"1'8, reddish. Fine field.  
 (6) Wispy, with two nuclei, enclosing faint pair.  
 (7) Has been thought variable.  
 (8) Moving together through space.  
 (9) In a fine scattered group.  
 (10) Struve calls it 'coloris egregie rubicundi, pæne rosei'; several pairs near.  
 (11) Probably binary, and common proper motion.  
 (12) Contains a red star, 8'5 mag. in the southern portion.

# MARCH (Mean Time of Transit at Greenwich).

## MARCH

7

NAME OF OBJECT	Visible	R.A.		Decl.	1			11			21			31			Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h. m.	° ' "		h. m.	° ' "	h. m.	° ' "	h. m.	° ' "	h. m.	° ' "	h. m.	° ' "	h. m.	° ' "				
Sidereal time at 9 P.M.	—	—	—	—	7 38	—	8 18	7 52	8 14	7 35	6 55	—	—	—	—	—	—	—	—	
Cluster in Monoceros	Dec.—Apr.	7 9.1	—	-10 6	8 31	—	7 52	7 16	6 33	5 57	6 33	—	—	—	—	—	—	—	Large scattered cluster (1).	
δ Geminorum	Dec.—Apr.	7 13.3	+22 11	8 35	8 35	+22 11	7 56	7 16	6 37	7 12	6 37	7.1	205	3.5, 8.9	—	—	—	—	Yellow, turquoise.	
63 Geminorum	Dec.—Apr.	7 20.9	+21 41	8 43	8 43	+21 41	8 3	7 24	6 45	7 24	6 45	44.6	324	5.5, 9.5	—	—	—	—	Orange, reddish purple (2).	
α Geminorum (Castor)	Dec.—Apr.	7 27.3	+32 8	8 49	8 49	+32 8	8 10	7 30	6 51	7 30	6 51	5.8	232	3.0, 4.0	—	—	—	—	Grisish, yellow, grnsh, yellow (2).	
Cluster in Gemini *	Dec.—Apr.	7 31.6	+21 51	8 53	8 53	+21 51	8 14	7 35	6 55	7 35	6 55	—	—	10.0-12.0	—	—	—	—	Faint, 12' diam.	
α Canis Minoris (Procyon)	Dec.—Apr.	7 33.3	+5 31	8 55	8 55	+5 31	8 16	7 36	6 57	7 36	6 57	35.0	80	1.0, 8.5	—	—	—	—	Yellowish white, blue (4).	
Cluster in Argo Navis	Jan.—Apr.	7 56.6	-14 33	8 58	8 58	-14 33	8 19	7 40	7 0	7 40	7 0	—	—	9.5	—	—	—	—	Visible to naked eye. 30' diam.	
2 Argūs	Jan.—Apr.	7 40.2	-14 24	9 2	9 2	-14 24	8 23	7 43	7 4	7 43	7 4	16.8	339	5.9, 6.7	—	—	—	—	Both white.	
Cluster in Argo Navis	Jan.—Apr.	8 5.3	-12 30	9 27	9 27	-12 30	8 48	8 8	7 29	8 8	7 29	—	—	5-10	—	—	—	—	20' by 12' (5).	
ζ Cancri	Jan.—Apr.	8 5.6	+18 0	9 27	9 27	+18 0	8 48	8 9	7 29	8 9	7 29	5.4	127	5.8, 6.5	—	—	—	—	White, yellow. Ternary (6).	
Pair in Lynx	Jan.—May	8 17.0	+42 22	9 39	9 39	+42 22	8 59	8 20	7 41	8 20	7 41	76.5	168	6.0, 8.0	—	—	—	—	Orange, blue.	
φ <sup>s</sup> Cancri	Jan.—Apr.	8 19.9	+27 19	9 41	9 41	+27 19	9 2	8 23	7 44	8 23	7 44	4.8	214	6.0, 6.3	—	—	—	—	Both white.	
72, 74 P. viii. Argūs	Jan.—Apr.	8 20.1	-23 40	9 42	9 42	-23 40	9 2	8 23	7 44	8 23	7 44	41.9	86	6.0, 8.5	—	—	—	—	Red, green.	
P. viii. 124-6 Cancri	Jan.—Apr.	8 33.3	+19 57	9 55	9 55	+19 57	9 16	8 36	7 57	8 36	7 57	92.8	61	6.3, 7.0	—	—	—	—	Deep yellow, white (7).	
Cluster in Cancer (Praesepe)	Jan.—Apr.	8 33.6	+20 20	9 55	9 55	+20 20	9 16	8 36	7 57	8 36	7 57	—	—	6.5-9	—	—	—	—	Contains about 50 large stars.	
ε Hydæ *	Jan.—Apr.	8 40.7	+6 50	10 2	10 2	+6 50	9 23	8 44	8 4	8 44	8 4	3.3	228	3.8, 7.4	—	—	—	—	Deep yellow, ashy blue. Binary.	
Red star in Cancer	Jan.—May	9 3.8	+31 26	10 25	10 25	+31 26	9 46	9 7	8 27	9 7	8 27	—	—	6.5	—	—	—	—	Pale red.	
38 Lynceis	Jan.—May	9 11.7	+37 17	10 33	10 33	+37 17	9 54	9 14	8 35	9 14	8 35	2.9	236	4.2, 6.3	—	—	—	—	White, ashy blue.	
21 Ursæ Maj.	Dec.—May	9 17.6	+54 30	10 39	10 39	+54 30	10 0	9 20	8 41	9 20	8 41	5.6	311	7.2, 8.2	—	—	—	—	White, bluish.	
α Hydæ	Jan.—May	9 22.0	-8 10	10 43	10 43	-8 10	10 4	9 25	8 45	9 25	8 45	281.3	153	2, 9.0	—	—	—	—	Pale red, pale green. A var.	
6 Leonis	Jan.—May	9 25.8	+10 13	10 47	10 47	+10 13	10 8	9 29	8 49	9 29	8 49	37.0	75	5.0, 9.5	—	—	—	—	Orange, deep blue.	
ο (14) Leonis	Jan.—May	9 35.1	+10 24	10 56	10 56	+10 24	10 17	9 38	8 58	9 38	8 58	80.9	41	3.8, 9.3	—	—	—	—	Reddish white, blue (8).	
9 Sextantis	Jan.—May	9 48.1	+5 29	11 9	11 9	+5 29	10 30	9 51	9 11	9 51	9 11	53.0	294	6.5, 8.5	—	—	—	—	Reddish, pale blue.	
Nebula in Sextans	Jan.—May	9 59.5	-7 10	11 21	11 21	-7 10	10 41	10 2	9 23	10 2	9 23	—	—	—	—	—	—	—	Milky white. 3' by 1'.	

- (1) A sapphire 6 mag. in the field.
- (2) Two other companions forming a curve 1.42' north is a pair 7.5, 9.0; 27.0°; 7.3; 14.0; 1.0/3; very white.
- (3) Binary, period very long, over 1,000 years. A 3rd star, 9.9, blue, at 156.0°; 5.8 mag. is a close pair, having a 6.6 magnitude star at 58.0; 1.0/0 (1866.0). 73.0/3 probably forms a ternary system; and a 4th, 9.5 mag., at 223.0; 210.0/0 may also be connected.
- (4) Distance of Procyon one million six hundred and eighty thousand times that of the Earth from the Sun. The 8.5 mag. is a very close double. Another
- star, 7.0 mag., 65.5" distant in the direction of 99.0, is a close pair; 7.0, 7.3; 14.0; 1.0/3; very white.
- (5) A bright orange 5.2 mag. s.f.
- (6) 5.8 mag. is a close pair, having a 6.6 magnitude star at 58.0; 1.0/0 (1866.0). The period of this pair is 61 years. C revolves round the common centre of gravity, but in an anomalous orbit.
- (7) A 7.3 mag. at 88.0; 99.0/8, white.
- (8) 3.8 mag. is moving.

## APRIL (Mean Time of Transit at Greenwich).

NAME OF OBJECT	Visible	R.A.		Decl.		1		11		21		30		Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h.	m.	°	'	h.	m.	h.	m.	h.	m.	h.	m.				
Sidereal time at 9 P.M.	—	—	—	—	—	9	41	10	20	10	59	11	35	—	0	—	—
Pair in Gemini.	Dec.—Apr.	6	2'3	+24	27	5	22	4	43	4	4	3	28	31'0	188	6'7, 8'2	Deep yellow, blue.
12 Lyonsis*	Dec.—May	6	36'2	+59	33	5	56	5	17	4	38	4	2	1'7	126	5'7, 6'4	Both white (1).
Pair in Auriga.	Dec.—May	6	36'3	+40	45	5	56	5	17	4	38	4	2	28'5	131	6'2, 8'4	Orange, turquoise.
μ Canis Majoris.	Dec.—Mar.	6	50'9	-13	54	6	11	5	32	4	52	4	17	2'9	339	5'2, 8'2	Yellow, turquoise.
Pair in Auriga.	Dec.—May	6	51'7	+37	15	6	12	5	32	4	53	4	18	25'9	56	6'8, 7'1	Both white.
Cluster in Canis Maj.	Dec.—Apr.	6	54'3	-13	32	6	14	5	35	4	56	4	20	—	—	8-10'5	20' diam.
19 Lyonsis	Dec.—May	7	13'5	+55	30	6	33	5	54	5	15	4	39	14'7	314	5'2, 6'2	Whitish blue, blue (2).
20 Lyonsis	Dec.—May	7	13'5	+50	22	6	33	5	54	5	15	4	39	15'0	253	6'9, 7'2	Both white.
Pair in Gemini.	Dec.—Apr.	7	25'9	+23	8	6	46	6	6	5	27	4	52	11'6	186	6'3, 8'2	White, blue.
α Geminorum (Castor)	Dec.—Apr.	7	27'3	+32	8	6	47	6	8	5	28	4	53	5'8	232	3'0, 4'0	Both grmish, yellow. Binary (3).
η Argūs	Dec.—Mar.	7	29'5	-23	14	6	49	6	10	5	31	4	55	9'0	288	6'0, 6'0	Red, blue.
Cluster in Argo.	Dec.—Apr.	7	31'3	-14	14	6	51	6	12	5	33	4	57	—	—	5'5-9'5	Visible to naked eye (4).
ξ Argūs	Dec.—Apr.	7	34'2	-26	33	6	54	6	15	5	35	5	8	10'0	318	5'0, 5'0	Both topaz yellow.
5 Argūs	Dec.—Apr.	7	42'6	-11	55	7	2	6	23	5	44	5	8	3'3	17	5'7, 7'5	White, olive.
Pair in Cancer.	Dec.—May	7	58'7	+27	52	7	18	6	39	6	0	5	24	3'3	353	6'5, 7'1	White, deep blue.
Cluster in Monoceros	Dec.—May	8	7'9	-5	27	7	28	6	48	6	9	5	34	—	—	8-10'5	20' diam.
Pair in Cancer.	Dec.—May	8	18'2	+20	32	7	38	6	59	6	19	5	44	37'5	191	7'0, 8'2	White, blue.
Pair in Monoceros.	Dec.—May	8	32'3	-6	24	7	52	7	13	6	33	5	58	64'6	211	6'0, 7'7	Yellow, whitish blue.
P. viii. 129 Caneri	Dec.—May	8	33'8	+20	5	7	53	7	14	6	35	5	59	20'0	55	5'7, 9'5	Deep yellow, blue (5).
31 Monocerotis	Dec.—May	8	38'0	-6	49	7	58	7	18	6	39	6	4	78'3	309	5'5, 7'5	Golden, deep blue.
P. viii. 160 Hydreæ	Dec.—May	8	39'6	-2	11	7	59	7	20	6	41	6	5	4'5	260	6'7, 7'7	White, ashy olive.
Caneri	Dec.—May	8	39'7	+29	11	7	59	7	20	6	41	6	5	30'4	307	4'2, 6'3	Deep yellow, deep blue.
Cluster in Cancer	Dec.—May	8	44'9	+12	14	8	4	7	25	6	46	6	10	—	—	9-10'5	About 200 stars, 20' diam. (6).
17 Hydreæ	Dec.—May	8	49'9	-7	32	8	9	7	30	6	51	6	15	4'2	358	6'7, 7'2	Both white.

(1) Binary; period perhaps 650 years. A 7'4 mag. at 307°; 8''7, bluish red.

(2) A blue 6'5 mag. at 358°; 2'15''2.

(3) Distance 1,950,000 times that of the Earth from the Sun (see March list).

(4) Contains several pairs.

(5) 7'0 mag., blue, at 342°; 65''4; and an 8'0 mag., also blue, at 44°; 82''5.

(6) Visible in the finder, shaped like a Phrygian cap.



APRIL

APRIL (Mean Time of Transit at Greenwich).

NAME OF OBJECT	Visible	R.A.		Decl.	1		11		21		30		Dis- tances	Posi- Angles	Mag.	COLOURS AND REMARKS
		h.	m.		h. m.	h. m.	h. m.	h. m.	h. m.	h. m.						
Sidereal time at 9 P. M.																
♂ Ursæ Maj.*	Dec.—June	9	0'4	+67 36	8 20	9 41	10 20	10 59	11 35	12 10	12 45	13 20	—	—	—	—
Pair in Cancer	Dec.—May	9	0'9	+23 26	8 20	7 41	7 41	7 2	6 26	6 26	6 26	6 26	2'4	233	5'1, 8'7	Yellow, reddish blue. Binary.
Nebula in Ursa Maj.	Dec.—June	9	14'1	+51 29	8 34	7 54	7 54	7 15	6 40	6 40	6 40	6 40	7'2	201	6'4, 7'1	White, ashy.
39 Lyncis	Dec.—June	9	14'8	+50 2	8 34	7 55	7 55	7 16	6 40	6 40	6 40	6 40	—	—	—	4' by 2'. White (1). White, blue.
41 Lyncis	Dec.—June	9	21'2	+46 7	8 41	8 1	8 1	7 22	6 47	6 47	6 47	6 47	82'5	162	5'7, 7'5	Deep yellow, bluish white.
γ Hydæ	Feb.—Apr.	9	23'4	- 2 16	8 43	8 4	8 4	7 24	6 49	6 49	6 49	6 49	64'9	3	5'5, 8'5	White, lilac (2).
Pair in Leo Minor	Feb.—June	9	28'2	+40 28	8 48	8 8	8 8	7 29	6 54	6 54	6 54	6 54	24'7	147	6'2, 7'2	Both yellowish white (3).
Nebula in Ursa Maj.	Dec.—June	9	46'0	+69 37	9 5	8 26	8 26	7 47	7 11	7 11	7 11	7 11	—	—	—	0' by 2½'. Nucleus = 8 m. star (4).
Pair in Ursa Maj.	—	10	8'7	+71 38	9 28	8 49	8 49	8 9	7 34	7 34	7 34	7 34	16'7	167	6'0, 7'0	White, blue.
ζ Leonis	Feb.—June	10	10'3	+23 59	9 30	8 50	8 50	8 11	7 36	7 36	7 36	7 36	318'7	342	3'4, 6'0	White, yellow (5).
γ Leonis	Feb.—June	10	13'7	+20 25	9 33	8 54	8 54	8 14	7 39	7 39	7 39	7 39	3'3	115	2'2, 3'4	Golden, greenish gold (6).
33 Sextantis	Mar.—June	10	37'4	+ 5 21	9 57	9 17	9 17	8 38	8 3	8 3	8 3	8 3	6'6	240	6'0, 7'0	Yellow, blue.
54 Leonis	Mar.—June	10	49'4	+25 21	10 9	9 29	9 29	8 50	8 15	8 15	8 15	8 15	6'3	105	4'8, 6'7	Pearly white, ashy green.
Scarlet star in Crater*	Mar.—June	10	55'0	-17 42	10 14	9 35	9 35	8 56	8 20	8 20	8 20	8 20	—	—	Var.	8'5-9'5. Intense blood col. (7).
α Ursæ Maj.	Dec.—June	10	56'7	+62 21	10 16	9 37	9 37	8 57	8 22	8 22	8 22	8 22	384'9	204	1'9, 8'0	Yellow, violet.
Nebula in Leo	Mar.—June	10	59'9	+ 0 35	10 19	9 40	9 40	9 1	8 25	8 25	8 25	8 25	—	—	—	3' by 1', wht. An 8 mg. star s.f.
Nebula in Ursa Maj.	Feb.—June	11	8'1	+55 38	10 27	9 48	9 48	9 9	8 33	8 33	8 33	8 33	—	—	—	Planet-like, pale, 2' diam.
ξ Ursæ Maj.*	Feb.—June	11	12'1	+32 10	10 31	9 52	9 52	9 13	8 37	8 37	8 37	8 37	1'7	242	4'0, 4'4	White or yellow (8).
ι Leonis	Mar.—June	11	17'9	+11 9	10 37	9 58	9 58	9 18	8 43	8 43	8 43	8 43	2'8	63	4'6, 7'4	Yellow, ashy olive.
τ Leonis	Mar.—June	11	22'1	+ 3 29	10 41	10 2	10 2	9 23	8 47	8 47	8 47	8 47	92'7	173	5'0, 6'7	Golden, bright blue.
57 Ursæ Maj.*	Mar.—June	11	22'9	+39 58	10 42	10 3	10 3	9 23	8 48	8 48	8 48	8 48	5'5	4	5'5, 8'3	White, violet.
γ Crateris	Apr.—June	11	26'6	-17 38	10 46	10 6	10 6	9 27	8 52	8 52	8 52	8 52	9'0	29	5'5, 5'8	Yellow, violet (9).
90 Leonis	Mar.—June	11	28'8	+28 25	10 48	10 9	10 9	9 29	8 54	8 54	8 54	8 54	3'2	212	6'5, 7'5	White, bluish (10).
2 Comæ Bereniceis	Mar.—June	11	58'4	+22 6	11 17	10 38	10 38	9 59	9 23	9 23	9 23	9 23	3'8	240	5'8, 7'2	White, ashy.

(1) In field with 37 Ursa, 6 mag., α, f. it. A 7 mag. south.  
 (2) Moving together through space.  
 (3) A 7.5 mag. in 95° 1, 118' 2.  
 (4) Another nebula 46' south, 6' long by 50" broad; 'splendens stria luminis.'  
 Two pairs in field.  
 (5) A 3rd star at 306' : 240'.  
 (6) Binary, period 407 years. The finest pair in the northern sky.  
 (7) In the field with α Crateris, 42 seconds of time following it. A blue star lies 60' from the scarlet one in 269°, and another s.f.  
 (8) The earliest calculated binary, period 60 years and nine months.  
 (9) Common proper motion.  
 (10) An 8.5 mag., reddish, at 235° : 64' 5.

MAY (Mean Time of Transit at Greenwich).

NAME OF OBJECT	Visible	R.A.		Decl.	1		11		21		31		Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h. m.	o ' /		h. m.	o ' /	h. m.	o ' /	h. m.	o ' /						
Sidereal time at 9 P.M.	—	—	—	—	11 39	12 18	12 58	13 37	14 16	14 55	15 34	16 13	—	0	—	—
Pair in Camelopardus	—	8 8.1	+72 46		5 30	4 51	4 11	3 32	2 52	2 12	1 32	1 05	43.9	86	5.7, 9.0	Deep yellow, blue.
v <sup>1</sup> Cancrī	Dec.—May	8 19.8	+24 54		5 42	5 2	4 23	3 44	2 43	1 42	8 41	5.8	5.8	41	6.2, 7.1	Greenish white, olive.
P. viii. 108 Hydre	Dec.—May	8 29.8	+7 1		5 52	5 12	4 33	3 54	2 53	1 52	10.5	27	10.5	27	6.1, 7.1	White, reddish ash.
ε Cancrī	Jan.—May	8 33.9	+19 58		5 56	5 16	4 37	3 58	2 57	1 56	134.2	249	134.2	249	5.7, 7.0	Yellow, ashy yellow.
Cluster in Cancer	Jan.—May	8 44.9	+12 14		6 7	5 27	4 48	4 9	3 8	2 8	—	—	—	—	9-10.5	20' diam.; see April list.
Nebula in Leo Minor *	Jan.—May	8 45.5	+33 51		6 7	5 28	4 49	4 9	3 8	2 8	—	—	—	—	—	5' by 1'; faint. 4 stars surround it.
Pair in Hydra	Feb.—May	8 51.6	-17 0		6 13	5 34	4 55	4 15	3 35	2 54	2.4	182	2.4	182	7.0, 7.3	Both white.
66 Cancrī	Jan.—June	8 54.3	+32 42		6 16	5 37	4 57	4 18	3 38	2 57	4.5	137	4.5	137	6.0, 8.2	White, bronze.
Pair in Ursa Maj.	Dec.—June	9 3.7	+62 8		6 25	5 46	5 7	4 27	3 46	2 65	24.9	26	24.9	26	6.5, 7.0	Both white.
40 Lyncis	Jan.—June	9 14.1	+34 52		6 36	5 56	5 17	4 38	3 57	3 16	202.3	33	202.3	33	3.1, 8.7	Orange, violet (1).
23 Ursæ Maj.	Dec.—June	9 22.6	+63 34		6 44	6 5	5 26	4 46	3 65	2 84	22.8	271	22.8	271	4.0, 9.0	White, deep blue.
Pair in Lynx	Dec.—June	9 23.4	+45 11		6 45	6 6	5 26	4 47	3 66	2 85	23.8	157	23.8	157	7.2, 8.6	Yellow, blue (2).
7 Leonis	Feb.—June	9 29.6	+14 53		6 51	6 12	5 33	4 53	4 12	3 31	42.6	80	42.6	80	6.5, 8.0	Bluish white, violet.
Pair in Leo Minor	Feb.—June	9 34.3	+39 29		6 56	6 17	5 37	4 58	4 17	3 36	3.4	283	3.4	283	6.8, 8.0	Deep yellow, deep blue.
Nebula in Ursa Maj.	—	9 39.4	+72 49		7 1	6 22	5 42	5 3	4 21	3 10	—	—	—	—	—	45" diam. Whitish. Good field.
Pair in Leo	Feb.—June	9 50.7	+20 18		7 12	6 33	5 54	5 14	4 33	3 52	30.4	175	30.4	175	6.6, 7.5	White, yellow.
α Leonis (Rigulus)	Feb.—June	10 2.3	+12 31		7 24	6 44	6 5	5 26	4 45	3 64	176.9	307	176.9	307	1.1, 8.3	Bluish white, olive (3).
γ Leonis	Feb.—June	10 13.7	+20 25		7 35	6 56	6 16	5 37	4 56	4 15	3.3	115	3.3	115	2.2, 3.4	Golden, greenish gold (4).
Nebula in Hydra	Mar.—June	10 19.3	-18 4		7 41	7 1	6 22	5 43	5 3	4 22	—	—	—	—	—	Pale blue, resembling Jupiter (5).
49 Leonis *	Feb.—June	10 29.0	+9 14		7 50	7 11	6 32	5 52	5 32	4 51	2.4	158	2.4	158	6.2, 8.4	White, turquoise.
Nebula in Leo	Feb.—June	10 37.9	+12 18		7 59	7 20	6 41	6 1	5 20	4 39	—	—	—	—	—	2' diam.
Pair in Hydra	Mar.—June	10 42.0	-14 40		8 3	7 24	6 45	6 5	5 14	4 23	70.5	23	70.5	23	6.7, 7.6	Both white (6).
Red star in Hydra	Mar.—June	10 46.1	-20 39		8 7	7 28	6 49	6 9	5 18	4 27	—	—	—	—	6.0	Fine red (7).
Pair in Ursa Maj.	Jan.—June	10 52.9	+59 31		8 14	7 35	6 56	6 16	5 35	4 54	34.5	38	34.5	38	6.5, 7.5	Yellow, blue.

(1) 8.7 has a minute companion at 18'.  
 (2) 7 mag. star in field, *z*.  
 (3) Moving together through space.  
 (4) See April list; α 7 mag. star at 20° : 239' has large proper motion; another, 83 mag., at 302° : 327".  
 (5) About 30" diam.  
 (6) 7.6 is double, with a companion of the same magnitude and colour at 18° : 6".  
 All three stars are nearly in the same straight line. 22' exactly south of this triple is P. x. 159 : 7.5, 8.5; 11.0 : 30".  
 (7) Has a blue 9.5 mag. in 90° : 30".

NAME OF OBJECT	Visible	R. A.		Decl.	Dis- tances				Mag.	COLOURS AND REMARKS
		h. m.	° ' "		31	21	11	1		
Sidereal time at 9 P.M.	—	h. m.	° ' "		h. m.	h. m.	h. m.	h. m.	—	—
Pair in Ursa Maj.	—	11 39	—		12 18	12 58	13 37	13 37	—	—
♄ Leonis	Mar.—June	11 79	+74 5		7 50	7 11	6 31	6 31	7 <sup>o</sup> , 7 <sup>5</sup>	Both white (1).
Pair in Crater	Mar.—June	11 10'9	— 3 2		7 53	7 14	6 34	6 34	288	Pale yellow, violet.
83 Leonis	Mar.—June	11 21'0	+ 3 38		8 3	7 24	6 44	6 44	97	White, lilac.
Pair in Ursa Maj.	Jan.—July	11 32'6	+45 44		8 14	7 35	6 56	6 56	150	White, deep blue.
93 Leonis	Mar.—June	11 42'1	+20 51		8 24	7 45	7 5	7 5	264	Yellowish white, ash (2).
Nebula in Ursa Maj.	Mar.—June	11 47'0	+37 37		8 29	7 50	7 10	7 10	355	Yellow, white (3).
Pair in Ursa Maj.	—	11 54'0	+71 17		8 36	7 57	7 17	7 17	—	90" diam.
Pair in Virgo	Mar.—June	12 3'6	-11 13		8 45	8 6	7 27	7 27	151	Both white.
2 Canum Venat.	April—July	12 10'4	+41 18		9 31	8 13	7 34	7 34	92	Yellow, olive (4).
Red star in Virgo	April—June	12 19'4	+ 1 25		9 40	8 22	7 43	7 43	259	Deep yellow, blue.
♁ Corvi	April—June	12 24'0	-15 53		9 45	8 26	7 47	7 47	7	Var.? Very full red.
Nebula in Virgo	April—June	12 26'2	+15 3		9 47	8 29	7 49	7 49	214	White, purple.
Pair in Corvus	April—June	12 35'3	-12 23		9 56	8 38	7 58	7 58	—	Spindle-shaped; 7' by 2' (5).
♄ Virginis	April—June	12 35'9	- 0 49		9 57	8 38	7 59	7 59	6'1, 6'2	Yellowish white (6).
♁ Canum Venat. (Cor. Caroli)	April—July	12 50'7	+38 56		10 12	8 53	8 14	8 14	156	Both yellow. Binary (7).
Pair in Draco	Mar.—Sep.	13 9'2	+67 55		10 30	9 12	8 32	8 32	228	Pale yellow, dusky yellow.
♄ Ursæ Maj. (Mizar)	Mar.—July	13 19'3	+55 31		10 40	9 22	8 42	8 42	179'0	Both yellowish white (8).
♄ Hydræ	May—June	13 30'5	-25 55		10 51	9 33	8 53	8 53	148	Greenish white, emerald (9).
81 Virginis	April—June	13 31'6	- 7 18		10 52	9 34	8 55	8 55	192	Deep yellow, yellow.
Pair in Virgo	April—June	13 31'9	+ 2 58		10 13	9 34	8 55	8 55	42	Both yellow.
Cluster in Canes Venat.	April—July	13 36'9	+28 57		10 18	9 39	9 0	9 0	32	Yellow, bluish olive.
Red star in Centaurus	May—June	13 42'6	-27 48		11 3	9 45	9 5	9 5	10-12	6' diam. Compressed mass of
7 Virginis	April—June	13 55'8	+ 2 6		11 17	9 58	9 19	9 19	6'2	Blood red (10).
									290	Yellow, light purple.

(1) The 7<sup>o</sup> mag. has a rapid proper motion and sensible parallax, distance 1,050,000 times that of the earth from the sun.  
 (2) A 9<sup>o</sup> mag. at 8<sup>h</sup>: 31<sup>m</sup>: 3<sup>s</sup>.  
 (3) A similar pair in field  $\alpha$ ,  $\delta$ : 7<sup>o</sup>, 8'5": 25<sup>s</sup>: 76<sup>h</sup>: 8.  
 (4) An 8<sup>o</sup> mag., blue, at 9<sup>h</sup>: 41<sup>m</sup>: 8<sup>s</sup>, moving.  
 (5) In the nebulous region of Virgo.  
 (6) A small star in 228°: 57<sup>m</sup>: 9<sup>s</sup>.  
 (7) Alternately variable. The period of revolution is about 180 years.  
 (8) An 8<sup>o</sup> mag., blue, at 235°: 120<sup>m</sup>: 9<sup>s</sup>.  
 (9) Alcor, 4'5 mag., forms a double star to the naked eye with Mizar. It is 707<sup>m</sup>: 2 distant on an angle of 72<sup>o</sup>: a 7'5 mag., the 'Sidus Ludovicianum, is at 101°: 510<sup>m</sup>: 4<sup>s</sup>.  
 (10) Has a blue companion in 109°: 78<sup>m</sup>: 0<sup>s</sup>.





NAME OF OBJECT	Visible	R.A.		Decl.	1		11		21		30		Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h.	m.		h. m.	o	h. m.	h. m.	h. m.	h. m.	h. m.	h. m.				
Sidereal time at 9 P.M.	—	—	—	—	13 41	14 20	15 0	15 35	15 0	15 35	15 0	15 35	—	—	—	—
♄ Boötis	Apr.—July	14 12.1	+51 53	—	9 31	8 52	8 12	7 37	8 12	7 37	8 12	7 37	38.2	33	4.6, 8.0	White, deep purple (1).
♂ Boötis	May—July	14 35.4	+16 54	—	9 54	9 15	8 36	8 0	8 36	8 0	8 36	8 0	5.9	103	4.1, 6.0	White, bright ash blue.
54 Hydræ	May—June	14 39.4	-24 57	—	9 58	9 19	8 40	8 4	8 40	8 4	8 40	8 4	8.9	129	5.0, 7.0	Deep yellow, reddish violet (2).
♄ Boötis	May—July	14 40.0	+27 33	—	9 59	9 19	8 40	8 5	8 40	8 5	8 40	8 5	3.0	329	2.5, 5.0	Deep yellow, fine blue (3).
♌ Libræ	May—June	14 44.6	-15 34	10 3	10 3	9 24	8 45	8 9	8 45	8 9	8 45	8 9	229.5	314	3.0, 5.5	Pale yellow, light grey (4).
♄ Boötis	May—July	14 46.1	+19 34	10 5	10 5	9 26	8 46	8 11	8 46	8 11	8 46	8 11	3.3	260	4.6, 6.7	Yellow, red. Binary (5).
P. xiv. 212 Libræ	May—June	14 50.8	-20 54	10 10	10 10	9 30	8 51	8 16	8 51	8 16	8 51	8 16	15.4	291	5.5, 7.5	Yellow, grey (6).
Pair in Virgo	May—July	14 58.5	+ 5 56	10 17	10 17	9 38	8 59	8 23	8 59	8 23	8 59	8 23	9.7	346	6.7, 7.0	Both white.
Pair in Boötis	May—July	15 7.7	+19 42	10 26	10 26	9 47	9 8	8 32	9 8	8 32	9 8	8 32	24.7	10	5.8, 6.8	Deep yellow, pearl.
♄ Boötis	May—Aug.	15 10.9	+33 44	10 29	10 29	9 50	9 11	8 36	9 11	8 36	9 11	8 36	105.2	79	3.4, 7.4	Yellow, blue (7).
Cluster in Libra	May—July	15 12.7	+ 2 31	10 31	10 31	9 52	9 13	8 37	9 13	8 37	9 13	8 37	—	—	9.5-11	Fine compressed cluster (8).
Pair in Corona Bor.	May—July	15 25.1	+25 54	10 44	10 44	10 4	9 25	8 50	9 25	8 50	9 25	8 50	3.1	93	6.7, 8.2	Golden, blue.
Pair in Libra	May—July	15 32.5	- 8 25	10 51	10 51	10 12	9 33	8 57	9 33	8 57	9 33	8 57	12.0	188	6.2, 6.3	White, greenish yellow.
♄ Coronæ Bor.	May—July	15 35.1	+ 37 0	10 54	10 54	10 14	9 35	9 0	9 35	9 0	9 35	9 0	6.2	302	4.4, 5.9	White, ash blue.
♄ Scorpionis *	May—July	15 58.1	-11 4	11 17	11 17	10 37	9 58	9 23	9 58	9 23	9 58	9 23	1.3	196	5.0, 5.4	White, pale yellow. Binary (9).
♄ Scorpionis	May—July	15 58.8	-19 30	11 17	11 17	10 38	9 59	9 27	9 59	9 27	9 59	9 27	13.4	25	2.7, 4.5	White, lilac (10).
♄ Herculis	May—July	16 2.9	+17 21	11 21	11 21	10 42	10 3	9 27	10 42	10 3	9 27	9 27	30.0	11	5.1, 6.1	Yellow, golden.
♄ Scorpionis *	May—July	16 5.5	-19 10	11 24	11 24	10 45	10 5	9 30	10 45	10 5	9 30	9 30	41.0	337	4.3, 7.0	White, bluish. Quadruplex (11).
49 Serpentis	May—July	16 8.0	+13 50	11 26	11 26	10 47	10 8	9 33	10 47	10 8	9 33	9 33	3.8	330	7.0, 7.1	Both white. Binary (12).
Cluster in Scorpio	May—July	16 10.3	-22 42	11 29	11 29	10 50	10 10	9 35	10 50	10 10	9 35	9 35	—	—	10-10.5	Very comprsd., like a comet (13).
♄ Coronæ Bor.	May—Aug.	16 10.4	+19 29	11 29	11 29	10 50	10 10	9 35	10 50	10 10	9 35	9 35	3.8	206	5.8, 6.7	White, blue. Binary (14).
♄ Herculis	May—July	16 16.9	+ 34 25	11 35	11 35	10 56	10 17	9 41	10 56	10 17	9 41	9 41	40.5	239	3.6, 8.8	Yellow, lilac.
♄ Scorpionis * (Antares)	May—July	16 22.4	-26 11	11 41	11 41	11 2	10 22	9 47	11 2	10 22	9 47	9 47	3.0	271	1.1, 7.3	Red, blue (15).
17 Draconis	Apr.—Aug.	16 33.5	+ 53 9	11 52	11 52	11 13	10 33	9 58	11 13	10 33	9 58	9 58	3.6	115	5.2, 6.3	White, ash green (16).

(10) A. 7 mag. at 31° : 520".  
 (11) 4.3 is a close pair, having a blue 6.7 mag. at 7° : 0".9, and the 7.0 at 41".0 is also double, with a bluish 8.0 mag. at 46° : 2".0.  
 (12) Large common proper motion. Long period binary.  
 (13) A temporary star, 6.5 mag., appeared in this cluster in 1860, and lasted about one month.  
 (14) Period very long, 850 years about. A faint star at 88° : 57".5.  
 (15) Both stars have the same small common proper motion.  
 (16) A 5.0 mag., 10 Draconis, at 194° : 50".3.

(1) Common proper motion.  
 (2) Common proper motion.  
 (3) Binary? i. Duplex pulcherrima."  
 (4) Common proper motion.  
 (5) Period 127 years.  
 (6) Pair resembling 61 Cygni, large common proper motion.  
 (7) Common proper motion.  
 (8) 42 seconds (of time) preceding and 19' north of 5 Serpentis (an unequal pair).  
 (9) Period 96 years, a 7.2 mag. bright violet, at 66° : 7".3. At 281". distance in the direction of 169° is a small pair, 7.3, 8.0 : 102' : 10".5; blue, with a faint star at 61".

NAME OF OBJECT	Visible	R.A.		Decl.		1		11		21		31		Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h. m.	° ' "	h. m.	° ' "	h. m.	° ' "	h. m.	° ' "	h. m.	° ' "	h. m.	° ' "				
Sidereal time at 9 P.M.	—	—	—	—	—	15 39	16 19	16 19	16 58	17 38	17 38	17 38	—	—	—	—	—
54 Virginis . . . . .	May—July	13 7.4	+18 13	6 28	5 49	5 10	4 31	5 2	5 10	4 31	5 2	5 2	6.3, 7.0	—	—	—	Yellowish, reddish.
34. 35 (Bode) Draconis . . . . .	—	13 9.2	+67 54	6 30	5 51	5 12	4 32	4 32	5 12	4 32	4 32	4 32	5.6, 6.0	—	—	—	Yellow, white (1).
Nebula in Canes Venat.*	May—Aug.	13 10.7	+42 38	6 32	5 52	5 13	4 34	5 13	5 13	4 34	4 34	4 34	—	—	—	—	7' by 3'. Faint.
1 Boötis . . . . .	Apr.—July	13 35.3	+20 32	6 56	6 17	5 38	4 58	6 17	5 38	4 58	4 58	4 58	6.2, 9.3	—	—	—	Blue, very blue (2).
83 Ursæ Maj. . . . .	May—Aug.	13 36.4	+55 16	6 57	6 18	5 39	4 59	6 18	5 39	4 59	4 59	4 59	5.0	—	—	—	Red orange (3).
P. xiii. 219, 220 Boötis . . . . .	Apr.—July	13 45.2	+21 50	7 6	6 27	5 48	5 8	6 27	5 48	5 8	5 8	5 8	6.3, 6.7	—	—	—	White, yellowish white (4).
κ Boötis . . . . .	May—Aug.	14 9.4	+52 19	7 30	6 51	6 12	5 32	6 51	6 12	5 32	5 32	5 32	4.5, 6.6	—	—	—	Lemon yellow, deep blue (5).
Pair in Boötēs . . . . .	Apr.—July	14 11.3	+20 40	7 32	6 53	6 14	5 34	6 53	6 14	5 34	4 1	176	6.4, 8.5	—	—	—	White, olive.
P. xiv. 69 Boötis . . . . .	Apr.—July	14 17.6	+8 5	7 39	6 59	6 20	5 41	6 59	6 20	5 41	6 2	188	5.7, 6.5	—	—	—	White, olive blue.
χ Turdi Solitarii . . . . .	May—July	14 19.1	+19 27	7 40	7 1	6 21	5 42	7 1	6 21	5 42	5 42	5 42	6.0, 7.2	—	—	—	Both white (6).
Pair in Boötēs . . . . .	Apr.—July	14 23.6	+28 49	7 44	7 5	6 26	5 47	7 5	6 26	5 47	25.5	262	6.1, 7.0	—	—	—	Both white.
ε Boötis . . . . .	Apr.—July	14 40.0	+27 33	8 1	7 22	6 42	6 3	7 22	6 42	6 3	3.0	329	2.5, 5.0	—	—	—	Deep yellow, fine blue.
μ Libræ* . . . . .	May—July	14 43.1	+13 41	8 4	7 25	6 45	6 6	7 25	6 45	6 6	1.4	335	5.2, 6.2	—	—	—	White, whitish blue.
39 Boötis . . . . .	Apr.—July	14 45.8	+49 11	8 7	7 27	6 48	6 9	7 27	6 48	6 9	3.5	45	5.5, 6.2	—	—	—	White, bluish white.
Pair in Boötēs'. . . . .	Apr.—July	14 56.2	+54 18	8 17	7 38	6 58	6 19	7 38	6 58	6 19	40.3	342	6.2, 7.0	—	—	—	Both white (7).
Pair in Boötēs . . . . .	Apr.—July	14 56.7	+47 43	8 17	7 38	6 59	6 20	7 38	6 59	6 20	35.5	157	6.0, 8.2	—	—	—	White purple.
44 Boötis . . . . .	Apr.—July	15 0.1	+48 6	8 21	7 42	7 2	6 23	7 42	7 2	6 23	5.0	241	5.0, 5.9	—	—	—	Yellow, ash green (8).
P. xv. 39 Boötis . . . . .	Apr.—July	15 12.4	+51 22	8 33	7 54	7 15	6 45	7 54	7 15	6 45	75.8	107	6.7, 8.5	—	—	—	Deep yellow, blue.
Pair in Serpens . . . . .	May—July	15 13.3	+10 59	8 34	7 55	7 15	6 36	7 55	7 15	6 36	13.2	172	6.3, 7.3	—	—	—	Both white.
μ Boötis . . . . .	May—July	15 20.2	+37 45	8 41	8 2	7 22	6 43	8 2	7 22	6 43	108.3	172	4.5, 6.5	—	—	—	Yellow, bluish yellow (9).
δ Serpentis . . . . .	May—July	15 29.4	+10 55	8 50	8 11	7 31	6 52	8 11	7 31	6 52	3.4	188	3.9, 5.5	—	—	—	Greenish yellow, ash green.
Pair in Corona Bor. . . . .	May—July	15 42.2	+36 48	9 3	8 24	7 44	7 44	8 24	7 44	7 44	5	307	7.0, 7.9	—	—	—	White, deep violet.
P. xv. 220 Serpentis . . . . .	May—July	15 51.5	+3 44	9 12	8 33	7 54	7 14	8 33	7 54	7 14	10.4	323	6.9, 8.4	—	—	—	White, olive.
ρ Coronæ Bor. . . . .	May—July	15 56.7	+33 39	9 17	8 38	7 59	7 19	8 38	7 59	7 19	76.1	92	6.0, 8.5	—	—	—	White, blue (10).

(1) An 8<sup>o</sup> mag; deep blue, at 232° : 120''/9.

(2) Two other blue stars in the field, one s. p., the other n. p.

(3) Brightened suddenly in August 1868.

(4) In field with 6 Boötis, a 5 mag. star.

(5) Moving together through space.

(6) 7<sup>a</sup> is a close pair, there being an 8<sup>a</sup> mag. at 102° : 1''/4.

(7) A small white nebula with a bright nucleus in the field, A.

(8) Colours and magnitudes variable. Binary, period 267 years.

(9) 6<sup>5</sup> is double with an 8 mag. companion at 100° : 0''/65; this forms a binary pair, period 270 years; the two form a ternary system with μ Boötis itself.(10) Change, owing to proper motion of 6<sup>o</sup> mag.

NAME OF OBJECT	Visible	R.A.		Decl.	1		11		21		31		Dis- tances	Posit Angles	Mag.	COLOURS AND REMARKS
		h	m.		h.	m.	h.	m.	h.	m.	h.	m.				
Sidereal time at 9 P.M.	—	—	—	—	15 39	16 19	16 19	16 19	16 58	17 38	17 38	17 38	—	0	—	—
Pair in Draco	—	16 5'2	+70 34	—	9 26	8 46	8 7	7 28	8 7	7 28	46'9	6 2, 8'2	84	—	—	Deep yellow, blue.
$\rho$ (5) Ophiuchi	June—July	16 18'7	-23 11	—	9 39	9 0	8 21	7 41	8 21	7 41	3 8	5'0, 5'3	357	—	—	Pale topaz, blue (1).
$\lambda$ Ophiuchi *	June—Aug.	16 25'2	+2 14	—	9 40	9 6	8 27	7 48	8 27	7 48	1 7	4'4, 5'4	44	—	—	White, ashy green. Binary (2).
Cluster in Hercules.	May—Sept.	16 37'6	+36 41	—	9 58	9 19	8 39	8 0	8 39	8 0	—	10-12	—	—	—	Superb. Visible to naked eye (3).
Cluster in Ophiuchus	June—Aug.	16 47'3	-1 45	—	10 2	9 22	8 43	8 4	8 43	8 4	—	9-11	—	—	—	Globular, 3' diam.
Pair in Draco	May—Oct.	16 47'2	+59 41	—	10 8	9 28	8 49	8 10	8 49	8 10	47'3	6'7, 7'7	350	—	—	Reddish yellow, blue.
Cluster in Ophiuchus	June—Aug.	16 51'2	-3 55	—	10 12	9 32	8 53	8 14	8 53	8 14	—	9-11	—	—	—	10' diam. (4).
Dracoenis.	May—Oct.	17 3'0	+54 37	—	10 23	9 44	9 5	8 25	9 5	8 25	2'4	5'0, 5'0	163	—	—	Rdsh. white, white. Binary (5).
$\beta$ Ophiuchi.	June—July	17 8'4	-26 25	—	10 29	9 50	9 10	8 31	9 10	8 31	4'3	5'5, 5'5	198	—	—	Reddish yellow, yellow (6).
$\alpha$ Herculis.	June—Sept.	17 9'4	+14 31	—	10 30	9 51	9 11	8 32	9 11	8 32	4'7	3'0, 5'5	116	—	—	Golden, perfect azure (7).
$\delta$ Herculis.	May—Sept.	17 10'3	+24 59	—	10 31	9 51	9 12	8 33	9 12	8 33	16'5	2'9, 8'2	186	—	—	Pale white, ruddy purple (8).
Cluster in Hercules*	May—Oct.	17 13'7	+43 15	—	10 34	9 55	9 15	8 36	9 15	8 36	—	10-11'5	—	—	—	Compressed cluster, faint.
$\epsilon$ Ophiuchi.	June—Aug.	17 38'8	+2 38	—	10 59	10 20	9 41	9 1	9 41	9 1	20'3	5'5, 6'2	94	—	—	Both white.
$\zeta$ Ophiuchi *	June—Aug.	17 56'9	-8 11	—	11 17	10 38	9 59	9 19	9 59	9 19	1 8	5'0, 6'0	256	—	—	White, ash. Binary (9).
$\eta$ Ophiuchi *	June—Aug.	17 59'7	+2 32	—	11 20	10 41	10 1	9 22	10 1	9 22	2'1	4'3, 6'2	20	—	—	Deep yel., rdsh. purp. Bin. (10).
Cluster in Scutum Sob.	July—Sept.	18 12'4	-13 50	—	11 33	10 53	10 14	9 35	10 14	9 35	—	9-10	—	—	—	Large, scatter'd, too stars at least.
Neb. in Scutum Sobieski	July—Sept.	18 13'0	-17 11	—	11 33	10 54	10 15	9 35	10 15	9 35	—	—	—	—	—	The 'Horse-shoe.' Vis. in finder
Cluster in Segittarius	July—Sept.	18 29'5	-23 59	—	11 50	11 10	10 31	9 52	11 10	10 31	9 52	—	—	—	—	Bright. Globular. [(11).
$\alpha$ Lyrae (Vega) *	June—Nov.	18 33'1	+38 41	—	11 53	11 14	10 35	9 55	10 35	9 55	49'2	0'2, 9'5	156	—	—	Pale sapphire, blue (12).
$\epsilon^1$ Lyrae	June—Nov.	18 40'6	+39 33	—	12 1	11 21	10 42	10 3	11 21	10 42	10 3	4'6, 6'3	16	—	—	White, blue (13).
$\epsilon^2$ Lyrae	June—Nov.	18 40'6	+39 30	—	12 1	11 21	10 42	10 3	11 21	10 42	10 3	5'1, 5'2	135	—	—	Both white.
Cluster in Antinous	June—Sept.	18 45'0	-6 25	—	12 5	11 26	10 47	10 7	11 26	10 47	10 7	8-10'5	—	—	—	Magnificent; fan-shaped; 12'
$\beta$ Lyrae	June—Nov.	18 45'9	+33 14	—	12 6	11 27	10 47	10 8	11 27	10 47	10 8	45'6	149	—	—	Whitish yellow, blue (14).
$\gamma$ Draconis.	May—Nov.	18 49'5	+59 15	—	12 10	11 30	10 51	10 12	11 30	10 51	31'4	4'7, 8'1	338	—	—	Bright yellow, blue.

(1) Two 8 mag. stars at 152'' and 161'' and a faint one at twice the distance. Back-ground otherwise black.

(2) Period 122 years, or perhaps nearly twice that.

(3) Small apertures only show a few stars sprinkled over it.

(4) 50'  $\beta$ , a little  $\eta$ , of 30 Oph., 5'0 mag., orange.

(5) Period 648 years.

(6) Binary, very large proper motion, in which a 7 mag. at 12' 10'' shares. A 9 mag. at 306' : 108'' from the close pair is not connected with this remarkable system. The two stars of the close pair are variable.

(7) Stationary, but common proper motion, 3'0 var. from 3'0 to 4'0 in about three months.

(8) Rectilinear motion.

(9) Period 218 years;  $\alpha$  9'5 mag. at 127' : 100''/3.

(10) Period 944 years. Distance of 70 Oph. 1,270,000 times that of the Earth from the Sun; mass of the system three times that of the Solar system; distance between the two stars equal to that of the planet Neptune from the Sun.

(11) In a magnificent group of stars. In a state of change.

(12) Distance of Vega 1,140,000 times that of the Earth from the Sun. Another star, variable, at 40' : 150''/4.

(13)  $\epsilon^1$ - $\epsilon^2$ , 173' : 207''/5. Several stars between.

(14)  $\beta$  Lyrae in 12' 3'5 to 4'5 mag. in 12d 21h 47m with double maximum and minimum. A 9'2 mag., yellow, at 318' : 66''/0, and a 9'4, lilac, at 19' : 85''/8. A minute pair s.f.

## AUGUST (Mean Time of Transit at Greenwich).

NAME OF OBJECT	Visible	R.A.		Decl.		1		11		21		31		Dis- tances	Posit. Angles	Mags.	COLOURS AND REMARKS
		h. m.	° ' "	h. m.	° ' "	h. m.	° ' "	h. m.	° ' "	h. m.	° ' "	h. m.	° ' "				
Sidereal time at 9 P.M.	—	—	—	—	—	17 42	18 21	19 9	19 40	19 40	19 40	19 40	19 40	—	0	—	—
Pair in <i>Hercules</i>	June—Sept.	16 07	+13 38	7 19	6 40	6 40	6 40	6 1	5 21	338	327	327	327	6.2, 7.3	—	Yellow, white.	
$\nu^2$ <i>Coronæ Bor.</i>	May—Sept.	16 18	+33 58	7 37	6 58	6 58	6 58	6 18	5 39	399.4	165	165	165	5.1, 5.6	—	Yellow, yellow (1).	
Cluster in <i>Ophiuchus</i> *	June—Aug.	16 26.1	-12 47	7 45	7 5	6 26	5 47	6 26	5 47	—	—	—	—	10-11	—	Faint, large.	
36, 37 <i>Herculis</i>	June—Sept.	16 34.9	+ 4 26	7 54	7 14	6 35	5 56	6 35	5 56	69.5	230	230	230	5.7, 6.7	—	White, blue.	
<i>Nebula in Hercules.</i>	June—Sept.	16 39.7	+24 1	7 58	7 19	6 40	6 0	6 40	6 0	—	—	—	—	—	—	Planetary, 8" diam., blue (2).	
43 <i>Herculis</i>	June—Sept.	16 40.3	+ 8 47	7 59	7 20	6 40	6 1	6 40	6 1	82.3	230	230	230	5.5, 9.0	—	Rose, blue.	
19 <i>Ophiuchi</i>	June—Sept.	16 41.4	+ 2 16	8 0	7 21	6 41	6 2	6 41	6 2	22.3	92	92	92	6.0, 9.0	—	White, ash.	
$\alpha$ 34 <i>Ophiuchi</i>	June—Sept.	16 58.9	+13 44	8 17	7 38	6 59	6 20	6 59	6 20	294.7	116	116	116	5.8, 6.3	—	White, reddish yellow (3).	
33 <i>Herculis</i>	June—Sept.	17 9.4	+14 31	8 28	7 49	7 9	6 30	7 9	6 30	4.7	116	116	116	3.0, 5.5	—	Golden, perfect azure (4).	
$\rho$ <i>Herculis</i>	June—Sept.	17 19.7	+37 15	8 38	7 59	7 20	6 40	7 20	6 40	3.8	313	313	313	4.6, 5.5	—	Greenish white, emerald (5).	
53 <i>Ophiuchi</i>	June—Sept.	17 29.2	+ 9 40	8 48	8 8	7 29	6 50	7 29	6 50	41.3	191	191	191	5.6, 7.3	—	White, reddish blue.	
$\nu^1$ <i>Draconis</i>	May—Oct.	17 29.9	+55 16	8 48	8 9	7 30	6 50	7 30	6 50	61.8	313	313	313	4.7, 4.7	—	Yellowish white, white (6).	
P. xvii. 200 <i>Herculis</i>	June—Sept.	17 36.4	+24 34	8 55	8 16	7 36	6 57	7 36	6 57	16.1	8	8	8	6.0, 8.5	—	Golden, turquoise.	
67 <i>Ophiuchi</i>	June—Sept.	17 54.9	+ 2 56	9 13	8 34	7 55	7 15	7 55	7 15	54.8	143	143	143	4.5, 7.7	—	White, deep blue.	
95 <i>Herculis</i>	June—Sept.	17 56.7	+21 36	9 15	8 36	7 56	7 17	7 56	7 17	6.0	261	261	261	5.0, 5.2	—	Green, red (7).	
Pair in <i>Hercules</i>	May—Oct.	18 0.1	+48 27	9 19	8 39	8 0	7 21	8 0	7 21	27.2	120	120	120	6.3, 8.0	—	White, sky blue.	
59 <i>Serpentis</i>	July—Sept.	18 21.4	+ 0 8	9 40	9 0	8 21	7 42	8 21	7 42	3.8	316	316	316	5.7, 7.7	—	Deep yellow, indigo blue.	
Cluster in <i>Serpens</i>	July—Sept.	18 22.0	+ 6 29	9 40	9 1	8 22	7 42	8 22	7 42	—	—	—	—	9-10	—	Visible to naked eye.	
39 <i>Draconis</i>	May—Oct.	18 22.2	+58 44	9 41	9 1	8 22	7 43	8 22	7 43	3.6	359	359	359	5.2, 8.1	—	White, blue (8).	
Pair in <i>Draco</i>	May—Oct.	18 31.3	+52 16	9 50	9 10	8 31	7 52	8 31	7 52	25.8	272	272	272	5.7, 8.0	—	Yellow, deep blue.	
$\zeta$ <i>Lyrae</i>	June—Nov.	18 40.8	+37 29	9 59	9 20	8 40	8 1	8 40	8 1	43.7	149	149	149	4.0, 6.0	—	White, green.	
Red star in <i>Scutum Sob.</i>	July—Oct.	18 44.0	- 8 2	10 2	9 23	8 44	8 4	8 44	8 4	—	—	—	—	7.5	—	Fine red.	
<i>Nebula in Lyra.</i>	June—Nov.	18 49.3	+32 53	10 8	9 28	8 49	8 10	8 49	8 10	—	—	—	—	—	—	Annular, 70" by 55".	
$\theta^{\theta}$ <i>Serpentis</i>	July—Sept.	18 50.5	+ 4 3	10 9	9 29	8 50	8 11	8 50	8 11	21.8	104	104	104	4.0, 4.4	—	Yellow, golden yellow (9).	

(1) A grey 6 mag.  $\nu^1$ ,  $\nu^2$ . Both  $\nu^1$  and  $\nu^2$  have small companions;  $\nu^1$  a 10.5 mag. at 23 $^{\circ}$ ;  $\nu^2$  a 9.5 mag., garnet, at 15 $^{\circ}$ ; 104 $^{\circ}$ .(2) Equal in light to 8 mag. star: a reddish 6 mag. in field  $s. f.$ (3) A 7 mag. at 13 $^{\circ}$ ; 238 $^{\circ}$ .

(4) See July list. One of the loveliest objects in the heavens.

(5) In field  $s. f.$  is a pair, 5.8, 8.5, 13 $^{\circ}$ ; 32 $^{\circ}$  6. Yellow, blue.

(6) Common proper motion.

(7) The colours have been thought variable.

(8) A 7.5 mag., ash blue, at 21 $^{\circ}$ ; 89 $^{\circ}$  3.

(9) Suspected of variability; fine field.

# AUGUST (Mean Time of Transit at Greenwich).

## AUGUST

NAME OF OBJECT	Visible	R.A.		Decl.	1		11		21		31		Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h. m.	°		h. m.	°	h. m.	°	h. m.	°	h. m.	°				
Sidereal time at 9 P.M.	—	—	—	—	18 59 <sup>o</sup>	— 4 12	10 17	9 38	18 21	19 10	19 40	—	—	—	—	—
15 Aquilæ . . . . .	July—Oct.	18 59 <sup>o</sup>	— 4 12	10 17	19 10	8 59	8 19	34.5	207	6 <sup>o</sup> , 7 <sup>5</sup>	—	—	—	—	White, red lilac.	
6 (Bode) Cygni . . . . .	May—Nov.	19 9 <sup>2</sup>	+ 49 40	10 28	9 48	9 9	9 30	9.7	219	5 <sup>8</sup> , 6 <sup>1</sup>	—	—	—	—	Deep yellow, yellow (1).	
η Lyrae . . . . .	June—Nov.	19 9 <sup>9</sup>	+ 38 57	10 28	9 49	9 9	8 30	28.2	84	4 <sup>1</sup> , 8 <sup>0</sup>	—	—	—	—	Bluish white, violet (2).	
θ Lyrae . . . . .	June—Nov.	19 12 <sup>4</sup>	+ 37 56	10 31	9 51	9 12	8 33	100.7	70	5 <sup>0</sup> , 8 <sup>0</sup>	—	—	—	—	Yellow, blue (3).	
β Cygni ( <i>Albireo</i> ) . . . . .	July—Nov.	19 26 <sup>1</sup>	+ 27 43	10 44	10 5	9 26	8 46	34.5	56	2 <sup>7</sup> , 5 <sup>3</sup>	—	—	—	—	Golden, blue (4).	
Red star in Sagittarius . . . . .	July—Sept.	19 27 <sup>8</sup>	— 16 37	10 46	10 7	9 27	8 48	—	—	6 <sup>5</sup>	—	—	—	—	Fine ruby.	
α Aquilæ ( <i>Altaïr</i> ) . . . . .	July—Oct.	19 45 <sup>2</sup>	+ 8 34	11 3	10 24	9 45	9 5	156.5	310	1 <sup>0</sup> , 9 <sup>0</sup>	—	—	—	—	White, violet.	
ε Draconis * . . . . .	—	19 48 <sup>6</sup>	+ 69 59	11 7	10 27	9 48	9 9	2.9	5	4 <sup>2</sup> , 7 <sup>2</sup>	—	—	—	—	Deep yellow, turquoise.	
Nebula in Vulpecula . . . . .	July—Nov.	19 54 <sup>6</sup>	+ 22 25	11 13	10 33	9 54	9 15	—	—	—	—	—	—	—	The 'umb-bell' Nebula.	
θ Sagittæ . . . . .	July—Nov.	20 4 <sup>9</sup>	+ 20 34	11 23	10 44	10 4	9 25	11.4	326	6 <sup>0</sup> , 8 <sup>1</sup>	—	—	—	—	White, turquoise (5).	
Red star in Capricornus . . . . .	Aug.—Sept.	20 10 <sup>4</sup>	— 21 40	11 28	10 49	10 10	9 30	—	—	Var.	—	—	—	—	6 <sup>5</sup> -9 <sup>0</sup> ? Pure ruby.	
α <sup>1</sup> Capricorni . . . . .	Aug.—Oct.	20 11 <sup>7</sup>	— 12 54	11 30	10 50	10 11	9 32	376.3	291	3 <sup>0</sup> , 4 <sup>0</sup>	—	—	—	—	Both deep yellow (6).	
σ Capricorni . . . . .	Aug.—Sept.	20 12 <sup>8</sup>	— 19 28	11 31	10 52	10 12	9 33	55.8	177	5 <sup>0</sup> , 8 <sup>3</sup>	—	—	—	—	Orange, violet.	
δ Capricorni . . . . .	Aug.—Sept.	20 23 <sup>4</sup>	— 18 57	11 41	11 2	10 23	9 43	22.1	240	6 <sup>0</sup> , 7 <sup>0</sup>	—	—	—	—	White, bluish white.	
γ Delphini . . . . .	July—Nov.	20 41 <sup>4</sup>	+ 15 43	11 59	11 20	10 41	10 1	11.3	271	4 <sup>0</sup> , 5 <sup>7</sup>	—	—	—	—	Reddish yellow, green (7).	
ε Equulei . . . . .	July—Oct.	20 53 <sup>4</sup>	+ 3 51	12 11	11 32	10 53	10 13	10.6	73	6 <sup>1</sup> , 7 <sup>4</sup>	—	—	—	—	White, ash blue (8).	
12 Aquarii . . . . .	July—Oct.	20 58 <sup>1</sup>	— 6 16	12 16	11 37	10 57	10 18	2.7	192	6 <sup>0</sup> , 8 <sup>0</sup>	—	—	—	—	Yellow, blue.	
61 Cygni . . . . .	Jan.—Nov.	21 1 <sup>8</sup>	+ 38 11	12 20	11 40	11 1	10 22	20.7	121	5 <sup>0</sup> , 6 <sup>0</sup>	—	—	—	—	Golden yellow, orange (9).	
P. xxi. 1 Cygni . . . . .	July—Nov.	21 3 <sup>8</sup>	+ 29 43	12 22	11 42	11 3	10 24	3.5	123	6 <sup>1</sup> , 8 <sup>1</sup>	—	—	—	—	White, blue (10).	
1 Pegasi . . . . .	July—Nov.	21 16 <sup>8</sup>	+ 19 19	12 35	11 55	11 16	10 37	36.2	311	4 <sup>3</sup> , 8 <sup>2</sup>	—	—	—	—	Deep yellow, lilac.	
Cluster in Pegasus . . . . .	July—Nov.	21 24 <sup>5</sup>	+ 11 39	12 42	12 3	11 24	10 44	—	—	10 5-11 5	—	—	—	—	Globular, 5' diam.	
Cluster in Aquarius . . . . .	July—Oct.	21 27 <sup>5</sup>	— 1 20	12 45	12 6	11 27	10 47	—	—	10 0-11 5	—	—	—	—	3 <sup>8</sup> diam.	
Cluster in Cygnus . . . . .	July—Nov.	21 28 <sup>1</sup>	+ 47 56	12 46	12 7	11 27	10 48	—	—	7-10	—	—	—	—	Splendid, 1 <sup>o</sup> diam.	
ε Pegasi . . . . .	July—Nov.	21 38 <sup>6</sup>	+ 9 21	12 56	12 17	11 38	10 58	140.4	322	2 <sup>4</sup> , 8 <sup>8</sup>	—	—	—	—	Deep yellow, violet.	

(1) Large common proper motion pair resembling 61 Cygni. Distance, 430,000 times that of the Earth from the Sun. Beautiful field.  
 (2) Three other pairs in large field.  
 (3) Fine field.  
 (4) One of the finest and brightest coloured pairs.  
 (5) A 6<sup>7</sup> mag., yellow, at 156<sup>o</sup>:154<sup>o</sup>:5 which is coarsely quadruple, and a very small double comes at 150<sup>o</sup>:154<sup>o</sup>:4. α has a 9 mag. lilac comes at 221<sup>o</sup>:44<sup>o</sup>:3, and several faint companions.  
 (6) α has a 9<sup>5</sup> mag. comes at 285<sup>o</sup>:1<sup>o</sup>:3. Binary.  
 (7) Motion probably rectilinear, not binary, as has been supposed, with very large common proper motion. One of the first stars whose distance was determined; recent measures give about 430,000 times the distance of the Earth from the Sun.  
 (8) 5<sup>7</sup> variable in colour. An 8 mag. at 141<sup>o</sup> 41'.  
 (9) A third, 10 mag., at 58<sup>o</sup>, a fourth more distant.

## SEPTEMBER (Mean Time of Transit at Greenwich).

NAME OF OBJECT	Visible	R.A.		Decl.	1		11		21		30		Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h.	m.		h. m.	h. m.	h. m.	h. m.	h. m.	h. m.						
Sidereal time at 9 P.M.	—	—	—	—	19 44	20 23	21 3	21 38	21 38	21 38	21 38	21 38	—	—	—	—
Pair in Ophiuchus .	June—Aug.	17 11	1 31	— 1	6 18	5 38	4 59	4 24	4 24	4 24	4 24	20'3	280	6.0, 8.7	White, bluish.	
Pair in Ophiuchus .	June—Sep.	17 22.5	+11 30	6 39	6 0	5 20	5 20	4 45	4 45	4 45	4 45	27'1	283	6.3, 7.3	White, blue.	
Pair in Hercules .	May—Sep.	17 45.6	+25 20	7 2	6 23	5 44	5 44	5 8	5 8	5 8	5 8	6'6	142	6.8, 8.4	White, ashy.	
Pair in Hercules .	May—Sep.	17 54.7	+30 3	7 11	6 32	5 53	5 53	5 17	5 17	5 17	5 17	19'5	278	6.9, 7.9	Reddish yellow, sky blue.	
Red star in Serpens .	June—Aug.	17 56.0	-12 19	7 13	6 33	5 54	5 54	5 19	5 19	5 19	5 19	—	—	6.8	Strong reddish.	
Nebula in Draco .	—	17 58.3	+66 38	7 15	6 36	5 56	5 56	5 21	5 21	5 21	5 21	—	—	—	Planetary. 20" diam. Blue.	
P. xvii. 362 Tauri Pon. .	July—Sep.	18 0.3	+12 0	7 17	6 38	5 58	5 58	5 23	5 23	5 23	5 23	6'6	258	6.3, 7.2	White, reddish olive.	
Pair in Draco .	May—Oct.	18 0.9	+56 26	7 17	6 38	5 59	5 59	5 23	5 23	5 23	5 23	38'4	24	6.8, 7.7	White, white (1).	
100 Hercules .	June—Oct.	18 3.2	+26 5	7 20	6 40	6 1	6 1	5 26	5 26	5 26	5 26	14'0	3	5.5, 5.5	Both greenish yellow.	
Pair in Draco .	June—Nov.	18 31.3	+52 15	7 48	7 8	6 29	6 29	5 54	5 54	5 54	5 54	25'8	272	5.7, 8.0	Deep yellow, deep blue.	
Pair in Taurus Pon. .	June—Sep.	18 39.9	+ 5 22	7 56	7 17	6 38	6 38	6 2	6 2	6 2	6 2	2'2	115	6.2, 6.6	Both white (2).	
5 Aquilæ .	June—Sep.	18 40.6	- 1 4	7 57	7 18	6 38	6 38	6 3	6 3	6 3	6 3	13'0	122	6.3, 7.3	White, clear blue (3).	
Pair in Lyra .	June—Nov.	18 50.7	+33 50	8 7	7 28	6 48	6 48	6 13	6 13	6 13	6 13	45'4	350	5.5, 7.6	Straw colour, purple (4).	
Pair in Aquila .	June—Sep.	18 54.7	+12 44	8 11	7 32	6 52	6 52	6 17	6 17	6 17	6 17	16'7	259	6.8, 8.2	Reddish orange, deep blue (5).	
Red star in Aquila .	June—Sep.	18 58.3	- 5 51	8 15	7 35	6 56	6 56	6 21	6 21	6 21	6 21	—	—	7.3	Var.? Fine deep red (6).	
Cluster in Lyra .*	June—Nov.	19 12.1	+29 59	8 28	7 49	7 10	7 10	6 34	6 34	6 34	6 34	—	—	10'5-11'5	A nebula in small telescopes.	
Pair in Vulpecula .	June—Nov.	19 15.5	+26 27	8 32	7 52	7 13	7 13	6 38	6 38	6 38	6 38	54'5	5	6.2, 6.3	Deep blue, red.	
2 Aquilæ .	July—Nov.	19 24.7	- 3 2	8 41	8 2	7 22	7 22	6 47	6 47	6 47	6 47	—	—	6.0	Fine orange red. Var.?	
Pair in Cygnus .	May—Nov.	19 31.3	+59 53	8 48	8 8	7 29	7 29	6 54	6 54	6 54	6 54	76'5	287	5.7, 7.2	Golden, blue.	
16 Cygni .	May—Nov.	19 38.8	+50 16	8 55	8 16	7 36	7 36	7 1	7 1	7 1	7 1	37'8	135	5.1, 5.2	Both dull yellow.	
1 Cygni .	June—Nov.	19 42.1	+33 28	8 58	8 19	7 40	7 40	7 4	7 4	7 4	7 4	25'7	72	4.8, 8.2	Deep yellow, reddish blue (7).	
5 Sagittæ .	June—Nov.	19 43.9	+18 51	9 0	8 21	7 42	7 42	7 6	7 6	7 6	7 6	8'6	313	5.2, 8.4	White, olive blue.	
57 Aquilæ .	July—Nov.	19 48.4	- 8 29	9 5	8 25	7 46	7 46	7 11	7 11	7 11	7 11	35'8	171	5.1, 6.0	Golden, blue (8).	
ψ Cygni .	May—Nov.	19 52.7	+52 8	9 9	8 30	7 50	7 50	7 15	7 15	7 15	7 15	3'4	183	5.0, 7.1	Whitish yellow, ashy blue.	

(1) 7.7 has an 8.2 mag. star at 14.0° : 5.0°.

(2) Slight motion in angle, distance constant at 3.0". Good test for small instruments.

(3) A faint star, which may be variable, at 14.6° : 27".5.

(4) 5.5 is a close and very unequal pair. All three moving together through space.

(5) In low power field with 11 Aquilæ,  $\mu, \nu$ . 11 Aq. is a delicate and beautiful pair. 5.1, 9.0 : 2.600 : 17.0", white, sky blue.(6)  $2^m 43^s$  12 Aquilæ, nearly on parallel.(7) Fine field. About 1.0"  $\mu, \nu$  south, is a little 8 mag. pair 3" apart. About 1.0"  $\nu, \zeta$  is the variable  $\nu$  Cygni.

(8) The colours should be watched; they are possibly variable.

# SEPTEMBER (Mean Time of Transit at Greenwich).

## SEPTEMBER

19

NAME OF OBJECT	Visible	R.A.		Decl.	1		11		21		30		Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h.	m.		h.	m.	h.	m.	h.	m.	h.	m.				
Sidereal time at 9 P.M.	—	—	—	—	19 44	20 23	21 3	21 38	21 38	21 38	21 38	21 38	—	—	—	—
Pair in Cygnus . . .	June—Nov.	19 59.5	+38 1	9 16	9 25	8 46	8 7	7 31	7 22	7 35	7 22	23 8	335	7.2, 8.4	—	Red, blue. Superb colours.
Pair in Aquila . . .	July—Oct.	20 9.1	-3 51	9 25	9 10	8 46	8 7	7 31	7 22	7 35	7 22	14.1	233	6.5, 7.7	—	White, olive blue.
6° $\alpha$ Cygni . . .	May—Nov.	20 10.0	+46 24	9 26	9 26	8 47	8 8	7 32	7 32	7 32	7 32	338.1	323	4.0, 5.5	—	Orange, cerulean blue (1).
$\beta$ Capricorni . . .	July—Oct.	20 14.6	-15 10	9 31	9 31	8 51	8 12	7 37	7 37	7 37	7 37	205.4	267	3.0, 6.5	—	Deep yellow, deep blue (2).
Pair in Cepheus . . .	May—Dec.	20 23.6	+56 16	9 40	9 40	9 0	8 21	7 46	7 46	7 46	7 46	26.3	118	6.0, 8.0	—	White, deep blue.
$\zeta$ Cygni . . .	July—Nov.	20 41.0	+30 17	9 57	9 57	9 18	8 38	8 3	8 3	8 3	8 3	6.5	60	4.4, 9.0	—	Orange, blue (3).
Pair in Cepheus . . .	May—Dec.	20 46.5	+58 19	10 3	9 23	8 44	8 3	8 3	8 3	8 3	8 3	115.2	262	6.5, 7.3	—	White, clear-red.
Cluster in Capricornus . . .	July—Nov.	20 47.1	-12 58	10 3	9 24	8 45	8 9	8 9	8 9	8 9	8 9	—	—	10.5-11.5	—	3' diam. Faint (4).
$\gamma$ Equulei . . .	July—Nov.	21 4.9	+9 38	10 21	9 42	9 2	8 27	362.0	8 27	8 27	8 27	362.0	153	4.3, 6.5	—	Pale yellow, bluish.
Cluster in Cygnus . . .	June—Dec.	21 25.3	+46 36	10 41	10 2	9 23	8 47	—	8 47	8 47	8 47	—	—	9.5-10.5	—	Large and coarse.
$\delta$ Cephei . . .	—	21 27.2	+70 4	10 43	10 4	9 25	8 49	13.5	8 49	8 49	8 49	13.5	250	3.5, 7.7	—	White, deep blue.
$\epsilon$ Capricorni . . .	July—Oct.	21 30.7	-19 58	10 47	10 7	9 28	8 53	47.5	8 53	8 53	8 53	47.5	69	4.5, 8.5	—	White, bluish.
Cluster in Capricornus . . .	July—Oct.	21 33.9	-23 41	10 50	10 11	9 31	8 56	—	8 56	8 56	8 56	—	—	10.5-11.0	—	Fairly bright, comet-like (5).
$\mu$ Cygni . . .	July—Nov.	21 39.0	+28 14	10 55	10 16	9 36	9 1	3.2	9 1	9 1	9 1	3.2	119	4.5, 6.3	—	Whitish yellow, blue (6).
Red star in Aquarius . . .	July—Nov.	21 40.6	-2 44	10 57	10 17	9 38	9 3	—	9 3	9 3	9 3	—	—	6.5	—	Very red. Var. in colour and [mag.?
Pair in Pegasus . . .	July—Nov.	21 48.9	+19 11	11 5	10 25	9 46	9 11	22.2	9 11	9 11	9 11	22.2	111	6.1, 8.1	—	Deep yellow, blue.
Red star in Cepheus . . .	July—Dec.	21 53.4	+63 4	11 9	10 30	9 51	9 15	—	9 15	9 15	9 15	—	—	5.4	—	Deep red orange.
$\delta$ Aquarii . . .	Aug.—Nov.	22 8.0	-21 38	11 24	10 45	10 30	9 30	4.8	9 30	9 30	9 30	4.8	115	6.8, 8.2	—	Topaz, blue (7).
P. xxii. 65 Lacertæ . . .	July—Nov.	22 13.9	+37 11	11 30	10 50	10 11	9 36	15.5	9 36	9 36	9 36	15.5	193	6.2, 8.3	—	White, deep blue.
$\zeta$ Aquarii . . .	Aug.—Nov.	22 23.0	-0 36	11 39	10 59	10 20	9 45	3.3	9 45	9 45	9 45	3.3	399	4.1, 4.2	—	Both whitish yellow (8).
8 Lacertæ . . .	July—Nov.	22 30.7	+39 3	11 46	11 7	10 28	9 52	22.5	9 52	9 52	9 52	22.5	186	6.0, 7.0	—	Whitish green, green (9).
$\gamma$ Aquarii . . .	Aug.—Nov.	22 41.6	-14 40	11 57	11 18	10 39	10 3	28.5	10 3	10 3	10 3	28.5	115	5.9, 9.2	—	White, amethyst.
Pair in Cepheus . . .	—	22 48.7	+67 23	12 4	11 25	10 46	10 10	69.1	10 10	10 10	10 10	69.1	281	5.7, 8.4	—	Opal, deep blue.
Pair in Cepheus . . .	July—Jan.	22 58.7	+59 49	12 14	11 35	10 56	10 20	33.9	10 20	10 20	10 20	33.9	276	6.2, 8.7	—	White, deep blue.

(1) A 7.0 mag, deep blue, at  $174^{\circ}$ ;  $1.07''$ . Fine field.  
 (2) A blue 8.5 mag, at  $134^{\circ}$ ;  $2.26''$ .  
 (3) In the midst of a wonderful branching nebulosity; invisible, however, with small instruments.  
 (4) Follows P. xx. 325. Capricorni, 6 mag.,  $2.2''$  exactly on parallel.  
 (5) Precedes  $\delta$  Capr. (5.8 mag.),  $1''$   $46''$ , and is 6". An 8 mag. star close *s.v.* the cluster.  
 (6) Common proper motion, and probably binary. A 7.0 mag., blue, at  $56^{\circ}$ ;  $208''$ , which does not belong to the system.  
 (7) An 8.5 mag. n. s.  
 (8) Binary of long period; 1.625 years.  
 (9) There is a faint star at  $28''$  from the 7.0 mag., and a yellow 9 mag. star at  $8.4''$  from the 6.0 mag.

## OCTOBER (Mean Time of Transit at Greenwich).

NAME OF OBJECT	Visible	R.A.		Decl.	1			11			21			31			Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h.	m.		h.	m.	h.	m.	h.	m.	h.	m.	h.	m.	h.	m.				
Sidereal time at 9 P.M.	—	—	—	—	21	42	22	22	23	3	23	40	—	—	—	—	—	—	—	
39 Draconis . . . . .	June—Nov.	18	22.2	+58 44	5	41	5	1	4	22	3	3	3	3	3	6	359	5.2, 8.1	White, blue (1).	
Red star in Lyra *	June—Nov.	18	28.3	+36 54	5	47	5	7	4	28	3	4	3	4	3	—	8	8	Intense crimson. Var.	
Pair in Lyra . . . . .	June—Nov.	18	39.5	+44 49	5	58	5	19	4	39	4	0	25.9	10	25.9	10	6.8, 8.0	6.8, 8.0	Reddish yellow, blue.	
ε Lyrae . . . . .	June—Nov.	18	40.6	+39 33	5	59	5	20	4	40	4	1	3.1	16	3.1	16	4.6, 6.3	4.6, 6.3	White, blue (2).	
ε <sup>e</sup> (5) Lyrae . . . . .	June—Nov.	18	40.6	+39 30	5	59	5	20	4	40	4	1	2.6	137	2.6	137	5.1, 5.2	5.1, 5.2	Both white.	
Pair in Draco . . . . .	June—Dec.	18	56.1	+62 15	6	15	5	35	4	56	4	17	16.9	124	16.9	124	6.2, 9.0	6.2, 9.0	Reddish yellow, blue.	
Pair in Lyra . . . . .	June—Nov.	19	4.2	+34 35	6	23	5	43	5	4	4	25	13.3	271	13.3	271	6.3, 8.2	6.3, 8.2	White, deep blue (3).	
Pair in Sagitta . . . . .	July—Nov.	19	15.8	+18 56	6	34	5	55	5	16	4	36	8.8	288	8.8	288	6.4, 8.3	6.4, 8.3	White, deep blue.	
Red star in Sagitta . . . . .	July—Nov.	19	21.3	+19 34	6	40	6	0	5	21	4	42	—	—	—	—	6.1	6.1	Orange (4).	
P. xix. 154 Cygni . . . . .	June—Nov.	19	23.7	+49 54	6	42	6	3	5	23	4	44	71.8	307	71.8	307	6.7, 7.7	6.7, 7.7	Yellow, deep blue.	
β Cygni (Albireo) . . . . .	July—Nov.	19	26.1	+27 43	6	44	6	11	5	26	4	46	34.5	56	34.5	56	2.7, 5.3	2.7, 5.3	Golden, blue (5).	
ε Sagittæ . . . . .	July—Nov.	19	32.1	+16 12	6	50	6	11	5	32	4	52	99.7	81	99.7	81	5.7, 7.8	5.7, 7.8	Golden, light blue.	
Cluster in Cygnus . . . . .	June—Nov.	19	37.3	+39 56	6	56	6	16	6	16	5	37	4.58	—	—	—	9.5-11.5	9.5-11.5	Fine; near Milky Way.	
Pair in Draco . . . . .	May—Nov.	19	38.3	+60 15	6	57	6	17	5	38	4	59	18.1	28	18.1	28	6.3, 8.0	6.3, 8.0	White, sky blue.	
P. xix. 276-7 Cygni . . . . .	June—Nov.	19	41.5	+35 49	7	0	6	21	5	41	5	2	14.7	126	14.7	126	6.2, 6.8	6.2, 6.8	Yellow, blue.	
P. xix. 278 Cygni . . . . .	June—Nov.	19	41.6	+34 44	7	0	6	21	5	41	5	2	38.1	28	38.1	28	6.0, 8.2	6.0, 8.2	Deep yellow, deep blue.	
26 Cygni . . . . .	May—Nov.	19	58.2	+49 47	7	16	6	37	5	58	5	18	41.7	146	41.7	146	5.3, 8.5	5.3, 8.5	Golden, whitish blue.	
Cluster in Vulpecula . . . . .	July—Nov.	20	7.2	+26 9	7	25	6	46	6	7	5	27	—	—	—	—	6.0-10.0	6.0-10.0	Large and rich.	
Pair in Cepheus . . . . .	May—Dec.	20	15.6	+55 2	7	34	6	54	6	15	5	36	2.9	341	2.9	341	6.0, 7.2	6.0, 7.2	White, blue.	
Pair in Cygnus . . . . .	June—Nov.	20	19.0	+42 37	7	37	6	58	6	19	5	39	96.4	63	96.4	63	6.5, 7.5	6.5, 7.5	Yellow, blue.	
Pair in Cygnus . . . . .	June—Nov.	20	34.3	+38 15	7	52	7	13	6	36	5	54	24.4	332	24.4	332	6.8, 8.5	6.8, 8.5	Yellow, blue (6).	
49 Cygni . . . . .	July—Nov.	20	36.4	+31 54	7	55	7	15	6	36	5	57	2.8	49	2.8	49	6.0, 8.2	6.0, 8.2	Yellow, blue.	
Pair in Cygnus . . . . .	June—Nov.	20	58.0	+39 3	8	16	7	37	6	57	6	18	18.8	300	18.8	300	6.5, 9.0	6.5, 9.0	Golden, emerald.	
P. xx. 465 Cygni . . . . .	June—Nov.	20	59.6	+41 10	8	18	7	38	6	59	6	20	57.4	185	57.4	185	5.8, 8.0	5.8, 8.0	White, deep blue.	

(4) In fine group of stars, one of which, 6.2 mag., is also of a fine red orange colour.  
 (5) Cf. August list. 5.3 has been thought variable. The colours of bright pairs are usually best shown by putting the stars *very* slightly out of focus, so as to enlarge the spurious disc.  
 (6) Rapid rectilinear motion.

(1) A 7.5 mag., ash blue, at 21° : 89'' .3.  
 (2) Cf. July list. A 10.0 mag. at 135° : 142'' .4 from ε<sup>1</sup> and 37° : 126'' .0 from ε<sup>2</sup>. Several other minute stars between the pairs, two of which, 45'' apart, form the 'duplex debilissima' of Struve.  
 (3) Another pair in field *m.f.* : 6.4, 7.4 : 260° : 17'' .1 ; yellow, olive blue.



OCTOBER (Mean Time of Transit at Greenwich).

OCTOBER

NAME OF OBJECT	Visible	R.A.		Decl.	1		11		21		31		Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h.	m.		h.	m.	h.	m.	h.	m.	h.	m.				
Sidereal time at 9 P.M.	—	—	—	—	h.	m.	h.	m.	h.	m.	h.	m.	—	—	—	—
δ Equulei *	July—Nov.	21	9'0	+ 9 33	8 27	7 48	7 8	6 29	40 0	21	4 5, 10'0	—	—	—	Yellow, purplish (1).	
3 Pegasi	July—Nov.	21	32'0	+ 6 7	8 50	8 11	7 31	6 52	39 2	349	5'0, 6'0	—	—	—	White, grey (2).	
P. xxi. 248 Cephei	June—Dec.	21	35'5	+56 58	8 53	8 14	7 35	6 56	11 8	121	5 8, 7'5	—	—	—	White, bluish (3).	
' Garnet' star in Cepheus,	June—Dec.	21	40'0	+58 15	8 58	8 19	7 39	7 0	—	—	Var.	—	—	—	Deep orange (4).	
Pair in Cepheus	June—Dec.	21	48'1	+55 16	9 6	8 27	7 47	7 8	19 5	195	5'4, 6'2	—	—	—	White, ashy green.	
Cluster in Cepheus	June—Dec.	21	54 8	+ 54 17	9 13	8 35	7 54	7 15	—	—	9-11	—	—	—	Very large, but poor.	
29 Aquarii	Aug.—Nov.	21	56'2	-17 31	9 14	8 35	7 55	7 16	3 7	244	7'0, 7'2	—	—	—	White, bluish.	
P. xxi. 401 Lacertæ	June—Dec.	22	0'3	+59 19	9 18	8 39	8 0	7 20	183 4	73	6 5, 7'5	—	—	—	Both white (5).	
Cluster in Lacerta.	July—Dec.	22	10 8	+49 18	9 29	8 49	8 10	7 31	—	—	8-10	—	—	—	Fine, but straggling (6).	
53 Aquarii	Aug.—Nov.	22	20'3	-17 19	9 38	8 59	8 20	7 40	8 2	305	6'0, 6'4	—	—	—	Both white.	
δ Cephei	July—Dec.	22	24'9	+57 50	9 43	9 3	8 24	7 45	40 8	192	Var. 5'7	—	—	—	Yellow, deep blue (7).	
μ Pegasi	Aug.—Dec.	22	37'7	+29 38	9 56	9 16	8 37	7 58	80 8	339	3'0, 9'0	—	—	—	Deep yellow, bluish.	
P. xxii. 219 Aquarii	Aug.—Nov.	22	42'0	- 4 49	10 0	9 20	8 41	8 2	3 6	253	6 8, 7'3	—	—	—	Yellow, white (8).	
γ Aquarii	Aug.—Nov.	22	43'6	-14 12	10 1	9 22	8 43	8 3	132 0	293	4'0, 9'0	—	—	—	Orange, blue.	
15 Lacertæ	Aug.—Dec.	22	46'9	+42 42	10 5	9 25	8 46	8 7	—	—	5'0	—	—	—	Fine orange red.	
83.84 Aquarii	Aug.—Nov.	22	59'3	- 8 19	10 27	9 38	8 58	8 19	261 0	147	5'5, 6'8	—	—	—	White, red.	
2 Cassiopeiæ	Aug.—Feb.	23	4'7	+58 43	10 22	9 43	9 4	8 24	167 9	163	6'1, 8'3	—	—	—	Yellowish white, bluish.	
ψ Aquarii	Aug.—Dec.	23	9'9	- 9 43	10 28	9 48	9 9	8 30	49 4	312	4'0, 8'5	—	—	—	Orange, sky blue (9).	
P. xxiii. 69 Aquarii	Aug.—Dec.	23	17 8	- 9 5	10 35	9 56	9 17	8 38	4 8	253	6 8, 7'7	—	—	—	Reddish white, greyish blue.	
Red star in Andromeda.	Aug.—Feb.	23	18'7	+40 58	10 36	9 57	9 18	8 38	—	—	6'5	—	—	—	Copper red.	
Nebula in Andromeda.	Aug.—Feb.	23	20 4	+41 54	10 38	9 59	9 19	8 40	—	—	—	—	—	—	Planetary, blue.	
δ Sculptoris	Oct.	23	43'0	-28 46	11 1	10 21	9 42	9 3	74 3	297	5'0, 9'0	—	—	—	White, bluish white.	
6 Cassiopeiæ *	Aug.—Feb.	23	43 2	+ 61 35	11 1	10 22	9 42	9 3	1 6	196	5'1, 7'6	—	—	—	Golden, turquoise.	
Pair in Cassiopeiæ.	Aug.—Feb.	23	55'5	+59 43	11 13	10 34	9 54	9 15	58 9	270	Var. 7'7	—	—	—	Ruby, deep blue (10).	

(1) δ is an excessively close pair, having the shortest period known among binary stars as yet, about 10<sup>1</sup>/<sub>2</sub> years.  
 (2) Pretty little pair δ south.  
 (3) A 7.5 mag., blue, at 340' : 19' 8.  
 (4) Sir W. Herschel's 'garnet' star, now orange. Variable, 4-6 mag., but period not well determined.  
 (5) A 9.5 mag. at 30' : 90' 3; a 6.7 mag. at 38' : 236' 7, and a distant 7.5 mag. 16' in length, followed by splendid field.  
 (6) A is var. 3.5 to 4.5 mag. in 5<sup>d</sup> 8<sup>h</sup> 47<sup>m</sup>.  
 (7) A is var. 3.5 to 4.5 mag. in 5<sup>d</sup> 8<sup>h</sup> 47<sup>m</sup>.  
 (8) An 8 mag. at 146' : 48' 5.  
 (9) Cf. January list.  
 (10) Magnificent colours. A var. 5'0-8'0 mag. (?) Cf. January list.

NOVEMBER (Mean Time of Transit at Greenwich).

NAME OF OBJECT	Visible	R.A.		Decl.		1		11		21		30		Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h. m.	°	h. m.	°	h. m.	°	h. m.	°	h. m.	°	h. m.	°				
Sidereal time at 9 P.M.	—	—	—	—	—	23 44	0 24	1 3	1 3	1 39	—	—	—	—	—	—	—
Cluster in Cygnus	June—Nov.	20 19 0	+40 21	5 35	4 56	4 17	3 41	3 41	3 41	3 41	—	—	9-10.5	—	—	Fine group.	
Pair in Cygnus	June—Nov.	20 30.2	+32 7	5 46	5 7	4 28	4 28	4 28	4 28	3 52	28.5	23.7	6.1, 8.4	—	—	Yellow, deep blue.	
48 P. xx. 243 Cygni	July—Nov.	20 33.0	+31 9	5 49	5 10	4 31	3 55	3 55	3 55	3 55	179.1	175	5.6, 6.0	—	—	Both whitish yellow.	
Pair in Capricornus	July—Nov.	20 41.9	-18 37	5 58	5 19	4 39	4 4	4 39	4 4	4 4	15.6	294	6.0, 7.0	—	—	Both white.	
P. xx. 376 Equulei *	July—Nov.	20 50.0	+ 4 6	6 6	5 27	4 48	4 12	4 48	4 12	4 12	2.0	287	6.2, 7.9	—	—	Orange, purple.	
λ Equulei	July—Nov.	20 56.6	+ 6 44	6 13	5 33	4 54	4 19	4 54	4 19	4 19	2.5	226	6.5, 6.5	—	—	Both white.	
Nebula in Aquarius	July—Nov.	20 57.9	-11 49	6 14	5 35	4 55	4 20	4 55	4 20	4 20	—	—	—	—	—	Planetary, pale blue (1).	
Pair in Cygnus	July—Nov.	21 2.1	+33 40	6 18	5 39	5 0	4 24	5 0	4 24	4 24	8.0	225	7.4, 8.0	—	—	Both white (2).	
Pair in Cygnus	July—Dec.	21 6.5	+47 4	6 23	5 43	5 4	4 29	5 4	4 29	4 29	134.2	190	6.1, 6.9	—	—	White, deep yellow (3).	
Pair in Pegasus	July—Nov.	21 27.7	+20 13	6 44	6 4	5 25	4 50	5 25	4 50	4 50	2.9	331	7.2, 7.7	—	—	Both white.	
Pair in Cepheus	July—Mar.	21 32.2	+66 15	6 48	6 9	5 30	4 54	5 30	4 54	4 54	180.1	26	6.0, 6.1	—	—	Deep yellow, white.	
76 Cygni	July—Dec.	21 37.0	+40 17	6 53	6 14	5 34	4 59	5 34	4 59	4 59	64.3	230	5.5, 9.0	—	—	White, blue.	
Red star in Cygnus	July—Dec.	21 38.6	+37 30	6 55	6 15	5 36	5 1	5 36	5 1	5 1	—	—	7.8	—	—	Very fine ruby.	
Cluster in Cepheus	July—Mar.	21 43.2	+65 17	6 59	6 20	5 41	5 5	5 41	5 5	5 5	—	—	9.5-11.5	—	—	Loose cluster, 10' diam.	
100 Aquarii (Boote)	July—Nov.	21 48.7	- 3 51	7 5	6 25	5 46	5 11	5 46	5 11	5 11	20.8	185	6.2, 9.0	—	—	White, blue.	
ξ Cephei	July—Mar.	22 0.4	+64 4	7 16	6 37	5 58	5 22	5 58	5 22	5 22	6.5	283	4.9, 6.6	—	—	Bluish white, ash blue (4).	
Cluster in Lacerta	July—Dec.	22 0.8	+45 56	7 17	6 37	5 58	5 23	5 58	5 23	5 23	—	—	8.5-10.5	—	—	About 50 stars in field.	
P. xxii. 33 Pegasi *	July—Dec.	22 8.9	+16 38	7 25	6 46	6 6	5 31	6 6	5 31	5 31	10.5	355	6.5, 9.1	—	—	Golden, blue (5).	
Pair in Cepheus	—	22 10.8	+72 44	7 27	6 47	6 8	5 33	6 8	5 33	5 33	28.9	348	5.5, 8.0	—	—	Deep yellow, turquoise.	
33 Pegasi *	July—Dec.	22 18.2	+20 17	7 34	6 55	6 16	5 40	6 16	5 40	5 40	2.1	180	6.1, 8.7	—	—	White, red (6).	
Pair in Cepheus	July—Mar.	22 18.4	+66 8	7 34	6 55	6 16	5 40	6 16	5 40	5 40	4.2	95	6.6, 7.4	—	—	Reddish white, deep blue.	
ξ Aquarii	Aug.—Dec.	22 23.0	- 0 36	7 39	7 0	6 20	5 45	6 20	5 45	5 45	3.3	329	4.1, 4.2	—	—	Both whitish yellow (7).	
5 Lacertæ	July—Dec.	22 24.8	+47 7	7 41	7 1	6 22	5 47	6 22	5 47	5 47	—	—	4.7	—	—	Red orange.	
Pair in Lacerta	July—Dec.	22 38.6	+38 53	7 55	7 15	6 36	6 0	6 36	6 0	6 0	2.8	280	6.2, 8.5	—	—	Golden, ash.	

(1) 25" by 17". Beas magnifying like a planet.  
 (2) Rectilinear motion.  
 (3) 6.1 is a close and unequal pair. 3<sup>m</sup> 51<sup>s</sup> f. 65 Cygni.  
 (4) Probably slow binary.  
 (5) Rectilinear motion.  
 (6) Common proper motion; a third star, 8.5 mag. blue, at 329° : 65" 6, is fixed in space.  
 (7) Binary; cf. September list.

NOVEMBER (Mean Time of Transit at Greenwich).

NAME OF OBJECT	Visible	R.A.		Decl.	1		11		21		30		Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h. m.	o /		h. m.	o /	h. m.	o /	h. m.	o /	h. m.	o /				
Sidereal time at 9 P.M.	—	23 20	—	23 44	8 18	7 39	6 59	6 24	8 5	146	6 6, 7 8	—	—	—	White, deep blue.	
P. xxii. 306 Pegasi	Aug.—Jan.	23 20	+ 32 12	23 44	8 18	7 39	6 59	6 24	8 5	146	6 6, 7 8	—	—	—	White, deep blue.	
Pair in Andromeda.	Aug.—Feb.	23 48	+ 47 21	23 44	8 21	7 41	7 2	6 27	15 0	253	6 7, 7 7	—	—	—	Yellowish white, rosy.	
94 Aquarii	Sep.—Dec.	23 13 1	- 14 5	23 13 1	8 29	7 50	7 10	6 35	13 6	346	5 0, 7 2	—	—	—	Deep yellow, deep blue (1).	
Pair in Pisces	Sep.—Jan.	23 18 4	+ 3 6	23 18 4	8 34	7 55	7 16	6 40	7 0	231	6 7, 8 9	—	—	—	Red, ashy blue.	
P. xxiii. 100-1 Cassiop.	Aug.—Feb.	23 24 7	+ 57 55	23 24 7	8 40	8 1	7 22	6 46	75 8	269	5 0, 7 0	—	—	—	Whitish blue, yellow (2).	
P. xxiii. 216-7 Pegasi	Sep.—Jan.	23 47 1	+ 11 17	23 47 1	9 3	8 23	7 44	7 9	18 7	282	6 3, 7 0	—	—	—	Yellow, ashy yellow.	
σ Cassiopeiæ	Aug.—Feb.	23 53 2	+ 54 59	23 53 2	9 9	8 30	7 50	7 15	2 9	327	4 8, 7 1	—	—	—	Greenish white, blue.	
Nebula in Pegasus	Sep.—Jan.	23 57 3	+ 15 30	23 57 3	9 13	8 34	7 54	7 19	—	—	—	—	—	—	Round disc, 75" diam.	
Pair in Andromeda	Aug.—Feb.	0 14 0	+ 37 36	0 14 0	9 30	8 50	8 11	7 36	64 0	18	7 3, 9 0	—	—	—	White, bluish (3).	
Cluster in Cassiopeiæ	—	0 20 7	+ 70 45	0 20 7	9 36	8 57	8 18	7 42	—	—	8-11	—	—	—	Large and straggling.	
55 Piscium	Sep.—Jan.	0 33 9	+ 20 49	0 33 9	9 49	9 10	8 31	7 55	6 4	193	5 2, 8 2	—	—	—	Yellow, turquoise.	
Great Neb. in Andromeda	Aug.—Feb.	0 36 2	+ 40 35	0 36 2	9 52	9 12	8 33	7 58	—	—	—	—	—	—	The 'Queen of the Nebulae' (4).	
Nebula in Cetus	Nov.	0 42 0	- 25 54	0 42 0	9 58	9 18	8 39	8 4	—	—	—	—	—	—	40' by 6', cream colour.	
P. o. 251 Piscium	Sep.—Feb.	0 53 5	+ 0 10	0 53 5	10 9	9 30	8 50	8 15	21 0	315	6 9, 8 0	—	—	—	Yellow, olive blue (5).	
Pair in Andromeda	Aug.—Feb.	0 53 6	+ 44 6	0 53 6	10 9	9 30	8 50	8 15	7 7	192	6 0, 7 0	—	—	—	White, ashy blue.	
77 Piscium	Sep.—Feb.	0 59 9	+ 4 18	0 59 9	10 15	9 36	8 57	8 21	32 8	83	6 1, 6 8	—	—	—	White, pale lilac.	
37 Ceti	Sep.—Feb.	1 8 7	- 8 13	1 8 7	10 24	9 45	9 6	8 30	49 5	331	5 5, 7 5	—	—	—	White, bluish white (6).	
P. i. 39-40 Cassiop.	July—Mar.	1 13 3	+ 64 4	1 13 3	10 29	9 49	9 10	8 35	52 5	350	6 2, 8 0	—	—	—	White, red.	
P. i. 85-87 Piscium	Oct.—Feb.	1 22 4	+ 7 22	1 22 4	10 38	9 59	9 19	8 44	69 5	99	6 0, 7 5	—	—	—	Yellow, blue.	
Pair in Andromeda.	Aug.—Mar.	1 42 1	+ 17 10	1 42 1	10 58	10 18	9 39	9 3	1 9	220	6 5, 7 2	—	—	—	White, blue (7).	
γ Arietis	Oct.—Mar.	1 47 2	+ 48 24	1 47 2	11 3	10 23	9 44	9 9	8 8	179	4 0, 4 2	—	—	—	Both white (8).	
ι Trianguli	Oct.—Mar.	2 5 7	+ 29 46	2 5 7	11 21	10 42	10 2	9 27	3 6	78	5 1, 6 7	—	—	—	Golden blue (8).	
P. ii. 38-39 Trianguli	Oct.—Mar.	2 10 8	+ 28 13	2 10 8	11 26	10 47	10 7	9 32	14 1	210	6 7, 7 5	—	—	—	Yellow, blue.	
γ Ceti	Oct.—Feb.	2 37 4	+ 2 45	2 37 4	11 53	11 13	10 34	9 59	2 8	290	3 4, 7 1	—	—	—	Whitish yellow, ash (8).	

(1) Moving together through space.  
 (2) Each of the stars is itself a close pair, and there are several other companions.  
 (3) Rapid rectilinear motion.  
 (4) See January and February lists.  
 (5) Moving, but probably not binary.  
 (6) Moving together through space.  
 (7) A blue 87 mag. at 179"; a yellow, violet, at 206".  
 (8) Cf. January list.

## DECEMBER (Mean Time of Transit at Greenwich).

NAME OF OBJECT	Visible	R.A.		Decl.	1		11		21		31		Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS
		h.	m.		h. m.	h. m.	h. m.	h. m.	h. m.	h. m.						
Sidereal time at 9 P.M.																
P. xxiii. 51 Androm.	—	23	14.2	+48 0	6 32	5 53	5 13	4 34	3 41	2 22	3 1	3 41	0	—	—	Both yellow.
Cluster in Cepheus	—	23	29.3	+72 18	6 47	6 8	5 28	4 49	3 55	2 22	3 1	3 41	78.9	305	8-11	Scattered.
19 Piscium	—	23	40.6	+ 2 51	6 58	6 19	5 40	5 0	4 49	3 55	2 22	3 41	—	—	6.2	Fine orange red.
P. xxiii. 223 Cassiop.	—	23	47.8	+50 53	7 6	6 26	5 47	5 8	4 24	3 55	2 22	3 41	42.4	197	6.0, 9.0	White, bluish white.
Cluster in Cassiop.	—	23	52.9	+59 23	7 11	6 31	5 52	5 13	4 24	3 55	2 22	3 41	—	—	7-9	Stars large.
Pair in Andromeda.	—	23	53.6	+33 6	7 11	6 32	5 53	5 13	4 24	3 55	2 22	3 41	3.0	206	6.0, 6.4	Both yellow (1).
Pair in Cassiop.	—	23	56.7	+65 28	7 14	6 35	5 56	5 16	4 24	3 55	2 22	3 41	15.2	70	5.8, 7.0	Golden, ultramarine.
α Andromedæ*	—	0	2.5	+28 28	7 20	6 41	6 2	5 22	4 24	3 55	2 22	3 41	71.0	273	2.0, 10.4	Bluish white, purplish (2).
Pair in Andromeda.	—	0	25.4	+32 57	7 43	7 4	6 24	5 45	4 24	3 55	2 22	3 41	56.3	85	5.3, 8.3	Reddish yellow, blue.
* Cassiopeidæ	—	0	26.5	+62 18	7 44	7 5	6 26	5 46	4 24	3 55	2 22	3 41	—	—	4.3	Surrounded by fine fields.
Pair in Andromeda.	—	0	29.0	+36 12	7 47	7 7	6 28	5 49	4 24	3 55	2 22	3 41	11.7	312	6.5, 8.6	Reddish gold, blue.
π Andromedæ	—	0	30.7	+33 5	7 48	7 9	6 30	5 50	4 24	3 55	2 22	3 41	36.3	173	4.2, 8.0	White, blue.
Cluster in Cassiop.	—	0	36.8	+61 10	7 54	7 15	6 36	5 56	4 24	3 55	2 22	3 41	—	—	8.5-9.5	Fine cluster.
P. o. 175-6 Androm.	—	0	40.2	+30 19	7 58	7 18	6 39	6 0	4 24	3 55	2 22	3 41	46.4	54	6.9, 7.0	Yellow, deeper yellow.
65 Piscium	—	0	43.7	+20 49	8 1	7 22	6 43	6 3	4 24	3 55	2 22	3 41	—	—	6.1, 6.4	Both whitish yellow.
26 Ceti	—	0	57.9	+ 0 46	8 15	7 36	6 57	6 18	4 24	3 55	2 22	3 41	16.0	252	6.2, 9.0	White, lilac.
ψ Piscium	—	0	59.5	+20 51	8 17	7 38	6 58	6 19	4 24	3 55	2 22	3 41	30.0	160	4.7, 4.9	Both greenish yellow (3).
σ Piscium	—	0	59.9	+31 34	8 17	7 38	6 59	6 20	4 24	3 55	2 22	3 41	56.0	204	6.2, 9.5	Deep yellow, blue (4).
Pair in Pisces	—	1	6.4	+31 28	8 24	7 45	7 5	6 26	4 24	3 55	2 22	3 41	19.6	249	6.7, 8.0	White, blue.
Pair in Cassiop.	—	1	29.8	+58 3	8 47	8 8	7 29	6 49	4 24	3 55	2 22	3 41	24.7	75	6.5, 7.7	Green, red.
Nebula in Perseus	—	1	35.1	+60 1	8 53	8 13	7 34	6 55	4 24	3 55	2 22	3 41	—	—	—	White, elliptical (5).
Red star in Cassiop.	—	1	47.4	+69 38	9 5	8 26	7 46	7 7	4 24	3 55	2 22	3 41	—	—	7.0	Fine red.
λ Arietis	—	1	51.6	+23 2	9 9	8 30	7 50	7 11	4 24	3 55	2 22	3 41	37.9	46	4.5, 7.0	White, olive blue.
α Piscium	—	1	56.1	+ 2 13	9 14	8 34	7 55	7 16	4 24	3 55	2 22	3 41	3.0	323	4.1, 5.4	Green, ash green (6).

(1) In slow movement.

(2) Changes, owing to proper motion of α.

(3) A 10 mag. star in 121° 98' 2".

(4) A 9.0 mag. at 23.6° 136' 4".

(5) Double nebula, line S. N. a little the brighter.

(6) Probably slow binary.

# DECEMBER

## DECEMBER (Mean Time of Transit at Greenwich).

NAME OF OBJECT	Visible	R.A.	Decl.	1				11				21				31				Dis- tances	Posit. Angles	Mag.	COLOURS AND REMARKS																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											
				h. m.	°	h. m.	°	h. m.	°	h. m.	°	h. m.	°	h. m.	°	h. m.	°	h. m.	°																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																															
Sidereal time at 9 P.M.	—	—	—	1 42	—	2 22	—	3 1	—	3 41	—	4 21	—	5 1	—	5 51	—	6 31	—	7 11	—	7 51	—	8 31	—	9 11	—	9 51	—	10 31	—	11 11	—	11 51	—	12 31	—	13 11	—	13 51	—	14 31	—	15 11	—	15 51	—	16 31	—	17 11	—	17 51	—	18 31	—	19 11	—	19 51	—	20 31	—	21 11	—	21 51	—	22 31	—	23 11	—	23 51	—	24 31	—	25 11	—	25 51	—	26 31	—	27 11	—	27 51	—	28 31	—	29 11	—	29 51	—	30 31	—	31 11	—	31 51	—	32 31	—	33 11	—	33 51	—	34 31	—	35 11	—	35 51	—	36 31	—	37 11	—	37 51	—	38 31	—	39 11	—	39 51	—	40 31	—	41 11	—	41 51	—	42 31	—	43 11	—	43 51	—	44 31	—	45 11	—	45 51	—	46 31	—	47 11	—	47 51	—	48 31	—	49 11	—	49 51	—	50 31	—	51 11	—	51 51	—	52 31	—	53 11	—	53 51	—	54 31	—	55 11	—	55 51	—	56 31	—	57 11	—	57 51	—	58 31	—	59 11	—	59 51	—	60 31	—	61 11	—	61 51	—	62 31	—	63 11	—	63 51	—	64 31	—	65 11	—	65 51	—	66 31	—	67 11	—	67 51	—	68 31	—	69 11	—	69 51	—	70 31	—	71 11	—	71 51	—	72 31	—	73 11	—	73 51	—	74 31	—	75 11	—	75 51	—	76 31	—	77 11	—	77 51	—	78 31	—	79 11	—	79 51	—	80 31	—	81 11	—	81 51	—	82 31	—	83 11	—	83 51	—	84 31	—	85 11	—	85 51	—	86 31	—	87 11	—	87 51	—	88 31	—	89 11	—	89 51	—	90 31	—	91 11	—	91 51	—	92 31	—	93 11	—	93 51	—	94 31	—	95 11	—	95 51	—	96 31	—	97 11	—	97 51	—	98 31	—	99 11	—	99 51	—	100 31	—	101 11	—	101 51	—	102 31	—	103 11	—	103 51	—	104 31	—	105 11	—	105 51	—	106 31	—	107 11	—	107 51	—	108 31	—	109 11	—	109 51	—	110 31	—	111 11	—	111 51	—	112 31	—	113 11	—	113 51	—	114 31	—	115 11	—	115 51	—	116 31	—	117 11	—	117 51	—	118 31	—	119 11	—	119 51	—	120 31	—	121 11	—	121 51	—	122 31	—	123 11	—	123 51	—	124 31	—	125 11	—	125 51	—	126 31	—	127 11	—	127 51	—	128 31	—	129 11	—	129 51	—	130 31	—	131 11	—	131 51	—	132 31	—	133 11	—	133 51	—	134 31	—	135 11	—	135 51	—	136 31	—	137 11	—	137 51	—	138 31	—	139 11	—	139 51	—	140 31	—	141 11	—	141 51	—	142 31	—	143 11	—	143 51	—	144 31	—	145 11	—	145 51	—	146 31	—	147 11	—	147 51	—	148 31	—	149 11	—	149 51	—	150 31	—	151 11	—	151 51	—	152 31	—	153 11	—	153 51	—	154 31	—	155 11	—	155 51	—	156 31	—	157 11	—	157 51	—	158 31	—	159 11	—	159 51	—	160 31	—	161 11	—	161 51	—	162 31	—	163 11	—	163 51	—	164 31	—	165 11	—	165 51	—	166 31	—	167 11	—	167 51	—	168 31	—	169 11	—	169 51	—	170 31	—	171 11	—	171 51	—	172 31	—	173 11	—	173 51	—	174 31	—	175 11	—	175 51	—	176 31	—	177 11	—	177 51	—	178 31	—	179 11	—	179 51	—	180 31	—	181 11	—	181 51	—	182 31	—	183 11	—	183 51	—	184 31	—	185 11	—	185 51	—	186 31	—	187 11	—	187 51	—	188 31	—	189 11	—	189 51	—	190 31	—	191 11	—	191 51	—	192 31	—	193 11	—	193 51	—	194 31	—	195 11	—	195 51	—	196 31	—	197 11	—	197 51	—	198 31	—	199 11	—	199 51	—	200 31	—	201 11	—	201 51	—	202 31	—	203 11	—	203 51	—	204 31	—	205 11	—	205 51	—	206 31	—	207 11	—	207 51	—	208 31	—	209 11	—	209 51	—	210 31	—	211 11	—	211 51	—	212 31	—	213 11	—	213 51	—	214 31	—	215 11	—	215 51	—	216 31	—	217 11	—	217 51	—	218 31	—	219 11	—	219 51	—	220 31	—	221 11	—	221 51	—	222 31	—	223 11	—	223 51	—	224 31	—	225 11	—	225 51	—	226 31	—	227 11	—	227 51	—	228 31	—	229 11	—	229 51	—	230 31	—	231 11	—	231 51	—	232 31	—	233 11	—	233 51	—	234 31	—	235 11	—	235 51	—	236 31	—	237 11	—	237 51	—	238 31	—	239 11	—	239 51	—	240 31	—	241 11	—	241 51	—	242 31	—	243 11	—	243 51	—	244 31	—	245 11	—	245 51	—	246 31	—	247 11	—	247 51	—	248 31	—	249 11	—	249 51	—	250 31	—	251 11	—	251 51	—	252 31	—	253 11	—	253 51	—	254 31	—	255 11	—	255 51	—	256 31	—	257 11	—	257 51	—	258 31	—	259 11	—	259 51	—	260 31	—	261 11	—	261 51	—	262 31	—	263 11	—	263 51	—	264 31	—	265 11	—	265 51	—	266 31	—	267 11	—	267 51	—	268 31	—	269 11	—	269 51	—	270 31	—	271 11	—	271 51	—	272 31	—	273 11	—	273 51	—	274 31	—	275 11	—	275 51	—	276 31	—	277 11	—	277 51	—	278 31	—	279 11	—	279 51	—	280 31	—	281 11	—	281 51	—	282 31	—	283 11	—	283 51	—	284 31	—	285 11	—	285 51	—	286 31	—	287 11	—	287 51	—	288 31	—	289 11	—	289 51	—	290 31	—	291 11	—	291 51	—	292 31	—	293 11	—	293 51	—	294 31	—	295 11	—	295 51	—	296 31	—	297 11	—	297 51	—	298 31	—	299 11	—	299 51	—	300 31	—	301 11	—	301 51	—	302 31	—	303 11	—	303 51	—	304 31	—	305 11	—	305 51	—	306 31	—	307 11	—	307 51	—	308 31	—	309 11	—	309 51	—	310 31	—	311 11	—	311 51	—	312 31	—	313 11	—	313 51	—	314 31	—	315 11	—	315 51	—	316 31	—	317 11	—	317 51	—	318 31	—	319 11	—	319 51	—	320 31	—	321 11	—	321 51	—	322 31	—	323 11	—	323 51	—	324 31	—	325 11	—	325 51	—	326 31	—	327 11	—	327 51	—	328 31	—	329 11	—	329 51	—	330 31	—	331 11	—	331 51	—	332 31	—	333 11	—	333 51	—	334 31	—	335 11	—	335 51	—	336 31	—	337 11	—	337 51	—	338 31	—	339 11	—	339 51	—	340 31	—	341 11	—	341 51	—	342 31	—	343 11	—	343 51	—	344 31	—	345 11	—	345 51	—	346 31	—	347 11	—	347 51	—	348 31	—	349 11	—	349 51	—	350 31	—	351 11	—	351 51	—	352 31	—	353 11	—	353 51	—	354 31	—	355 11	—	355 51	—	356 31	—	357 11	—	357 51	—	358 31	—	359 11	—	359 51	—	360 31	—	361 11	—	361 51	—	362 31	—	363 11	—	363 51	—	364 31	—	365 11	—	365 51	—	366 31	—	367 11	—	367 51	—	368 31	—	369 11	—	369 51	—	370 31	—	371 11	—	371 51	—	372 31	—	373 11	—	373 51	—	374 31	—	375 11	—	375 51	—	376 31	—	377 11	—	377 51	—	378 31	—	379 11	—	379 51	—	380 31	—	381 11	—	381 51	—	382 31	—	383 11	—	383 51	—	384 31	—	385 11	—	385 51	—	386 31	—	387 11	—	387 51	—	388 31	—	389 11	—	389 51	—	390 31	—	391 11	—	391 51	—	392 31	—	393 11	—	393 51	—	394 31	—	395 11	—	395 51	—	396 31	—	397 11	—	397 51	—	398 31	—	399 11	—	399 51	—	400 31	—	401 11	—	401 51	—	402 31	—	403 11	—	403 51	—	404 31	—	405 11	—	405 51	—	406 31	—	407 11	—	407 51	—	408 31	—	409 11	—	409 51	—	410 31	—	411 11	—	411 51	—	412 31	—	413 11	—	413 51	—	414 31	—	415 11	—	415 51	—	416 31	—	417 11	—	417 51	—	418 31	—	419 11	—	419 51	—	420 31	—	421 11	—	421 51	—	422

## CIRCUMPOLAR OBJECTS.

NAME OF OBJECT	R.A.	Decl.	Dis- tances	Posit. Angles	Mags.	COLOURS AND REMARKS
Pair in <b>Cepheus</b>	h. m.	° ' "	" "	102°	6'4, 7'1	Yellow, whitish yellow.
Cluster in <b>Cepheus</b>	0 7'7	+75 24	77"1	—	9-11'5	Very large, 150-200 stars.
α <b>Ursæ Min. (Polaris)</b>	1 17'0	+88 42	18'3	214	2'0, 8'4	Yellowish white, blue (1).
Pair in <b>Cepheus</b>	1 44'9	+75 39	3'2	247	7'0, 7'7	Both white.
Pair in <b>Cassiopeia</b>	1 52'9	+73 17	5'6	191	6'0, 9'0	White, blue.
47 <b>Cassiopeia</b>	1 53'7	+76 44	95"1	196	4'7, 9'2	White, blue.
Pair in <b>Cepheus</b>	1 59'7	+79 9	55"8	275	6'1, 6'8	Both white.
P. ii. 191 <b>Cephei</b> *	2 50'9	+78 57	4'5	230	5'4, 9'4	Golden, blue.
P. iv. 269 <b>Camelopardi</b>	5 3'5	+79 6	17'6	5	4'8, 8'3	Light yellow, blue (2).
Pair in <b>Camelopardus</b>	6 48'7	+75 24	12'0	28	7'0, 8'0	Yellow, deep blue.
Red star in <b>Camelop..</b>	7 7'1	+82 38	—	—	5'5	Light red.
Pair in <b>Camelopardus</b>	8 7'5	+72 46	43'9	86	5'7, 9'2	Yellow, blue.
Nebula in <b>Camelop..</b>	8 40'6	+78 39	—	—	—	Pretty bright, long.
Pair in <b>Ursa Maj.</b>	9 27'0	+73 36	4'9	134	6'8, 7'1	Both white.
Red star in <b>Camelop..</b>	11 54'4	+81 29	—	—	6'2	Fine orange.
Pair in <b>Camelopardus</b>	12 5'8	+82 21	65"1	76	6'0, 8'0	Orange, deep blue.
Pair in <b>Camelopardus</b>	12 11'2	+80 46	14'3	218	6'0, 7'0	Both white.
Pair in <b>Camelopardus</b>	12 48'1	+84 3	21'8	327	4'5, 5'0	Whitish yel., bluish white.
π' <b>Ursæ Min.</b>	15 36'1	+80 52	30'6	82	6'1, 7'0	Yellow, bluish white (3).
ψ <b>Draconis</b>	17 44'0	+72 12	30'9	15	4'1, 5'2	Yellow, lilac (4).
40, 41 <b>Draconis</b>	18 8'6	+79 59	20'3	235	5'4, 5'8	Both yellow (5).
Pair in <b>Draco</b>	18 57'4	+75 38	5'7	219	6'5, 7'2	Both white.
Red star in <b>Draco</b>	19 25'6	+76 20	—	—	6'5	Fine red. Var.?
κ <b>Cephei</b>	20 12'7	+77 22	7'4	123	4'5, 8'1	White, violet.
Pair in <b>Cepheus</b>	22 2'4	+82 19	13'6	75	6'1, 6'9	Yellow, bluish yellow.

(1) The companion was seen in sunlight by Struve with the Dorpat refractor of 9'6-in. aperture, February 9, 1826. Distance of Polaris two million two hundred and seventy thousand times greater than that of the Earth from the Sun.

(2) In rapid motion, possibly orbital?  
 (3) Moving together through space.  
 (4) Moving together through space.  
 (5) A 7 mag., lilac, in field, about 3½' s. f.

## DAILY ACCELERATION OF SIDEREAL ON MEAN TIME.

day	h. m.	days	h. m.	days	h. m.	days	h. m.	days	h. m.	days	h. m.	days	h. m.	days	h. m.
1	0 4	12	0 47	23	1 30	34	2 14	44	2 53	54	3 32	64	4 12	74	4 51
2	0 8	13	0 51	24	1 34	35	2 18	45	2 57	55	3 36	65	4 16	75	4 55
3	0 12	14	0 55	25	1 38	36	2 22	46	3 1	56	3 40	66	4 19	76	4 59
4	0 16	15	0 59	26	1 42	37	2 25	47	3 5	57	3 44	67	4 23	77	5 3
5	0 20	16	1 3	27	1 46	38	2 29	48	3 9	58	3 48	68	4 27	78	5 7
6	0 24	17	1 7	28	1 50	39	2 33	49	3 13	59	3 52	69	4 31	79	5 11
7	0 28	18	1 11	29	1 54	40	2 37	50	3 17	60	3 56	70	4 35	80	5 15
8	0 31	19	1 15	30	1 58	41	2 41	51	3 21	61	4 0	71	4 39	81	5 18
9	0 35	20	1 19	31	2 2	42	2 45	52	3 24	62	4 4	72	4 43	82	5 22
10	0 39	21	1 23	32	2 6	43	2 49	53	3 28	63	4 8	73	4 47	83	5 26
11	0 43	22	1 26	33	2 10									93	6 6

## OBJECTS SUITABLE FOR FOUR TO SEVEN-INCH TELESCOPES.

NAME OF OBJECT	Visible	R.A.	Decl.	Distances	Posit. Angles	Magnitude
34 Piscium . . .	Dec.—Feb.	h. m. 0 4'2	+10° 31'	" 7'8	160°	5'9, 10'2
26 Andromedæ . . .	Aug.—Feb.	0 12'7	+43 9	5'9	240	6'0, 10'1
42 Piscium . . .	Dec.—Feb.	0 16'5	+12 51	29'7	338	6'6, 11'0
12 Ceti . . .	Dec.—Feb.	0 24'2	- 4 35	8'6	187	6'0, 10'9
P. O. 245 Piscium . . .	Oct.—Feb.	0 52'3	+20 47	0'75*	93	6'7, 8'2
P. i. 11 Piscium . . .	Sept.—Feb.	1 6'7	+29 28	10'6	258	6'0, 10'0
φ Piscium . . .	Oct.—Feb.	1 7'5	+23 59	7'7	227	5'2, 10'0
42 Ceti (1) . . .	Dec.—Feb.	1 14'0	- 1 6	1'45	355	6'7, 7'5
ψ Cassiopeia (2) . . .	Aug.—Feb.	1 17'9	+67 32	28'7	107	4'5, 9'7
P. i. 123 Piscium (3) . . .	Nov.—Feb.	1 30'1	+ 7 4	1'33	33	7'2, 7'4
103 Piscium (4) . . .	Nov.—Feb.	1 33'1	+16 3	1'31	297	6'9, 9'0
ε Trianguli . . .	Nov.—Feb.	1 56'3	+32 44	4'0	118	5'6, 10'8
48 Cassiopeia . . .	July—Mar.	1 52'6	+70 21	0'96	267	5'0, 7'0
10 Arietis (5) . . .	Nov.—Feb.	1 57'1	+25 23	1'04	50	5'8, 8'0
P. ii. 96 Arietis . . .	Nov.—Mar.	2 23'9	+24 44	12'1	182	6'3, 10'7
γ Ceti . . .	Dec.—Mar.	2 29'9	+ 5 6	7'8	84	5'0, 9'9
Pair in Cetus . . .	Dec.—Mar.	2 35'0	+ 4 24	1'71	295	7'0, 8'9
84 Ceti (5) . . .	Dec.—Mar.	2 35'4	- 1 10	4'7	324	5'8, 9'6
θ Persei (6) . . .	Sept.—Mar.	2 36'4	+48 45	17'2	299	4'2, 9'8
Pair in Perseus . . .	Oct.—Mar.	2 40'1	+35 6	1'53	161	6'3, 8'7
π Arietis . . .	Dec.—Mar.	2 42'9	+17 0	3'3, 25'2	122, 109	5'5, 8'2, 11'0
41 Arietis . . .	Dec.—Mar.	2 43'4	+26 48	21'2, 34'0	266, 204	4'1, 11'3, 11'0
Pair in Cepheus . . .	—	2 50'9	+78 58	4'5	230	5'4, 9'4
ε Arietis (7) . . .	Dec.—Mar.	2 52'6	+20 54	1'39	201	5'4, 6'3
50 Arietis . . .	Dec.—Mar.	2 54'1	+17 33	2'1	67	7'1, 9'9
ρ <sup>2</sup> Eridani (8) . . .	Dec.—Feb.	2 57'1	- 8 8	2'5	85	5'4, 9'5
Pair in Cassiopeia (9) . . .	Aug.—Apr.	3 1'3	+71 7	0'98	213	7'0, 7'0
12 Eridani (10) . . .	Dec.—Jan.	3 7'2	-29 27	2'6	312	4'0, 9'5
τ <sup>4</sup> Eridani (11) . . .	Dec.—Jan.	3 14'5	-22 11	5'4	287	5'0, 9'5
Pair in Perseus . . .	Nov.—Mar.	3 17'3	+33 8	3'9	153	6'4, 9'9

(1) Binary.

(2) 9'7 is double, 10'6 : 256° : 3''1.

(3) Slow binary : a 10'0 mag. at 71° : 77''3.

(4) *n. β.*, 105 Piscium ; in the same field.

(5) Moving together through space.

(6) Moving together through space ; a 9'0 mag. at 218° : 69''2, not connected with the system of θ Persei.

(7) Binary.

(8) 9'5 variable.

(9) Binary, an 11'6 mag. in 307° : 20''.

(10) Sometimes called α Fornacis. Very rapid common proper motion.

(11) A 10'5 at 100° : 40''0 ; a 10'7 at 293° : 123''0 ; a 10'7 at 276° : 130''0 ; and a 10'0 at 236° : 160''1.

\* Distances under 2''0 are given in this list to the nearest hundredth of a second of arc.

NAME OF OBJECT	Visible	R. A.	Decl.	Distances	Posit. Angles	Magnitude
2 (Hevel) Camelopardi	Oct.—Mar.	h. m. 3 19'8	+59 32'	2"2	163 <sup>o</sup>	4'9, 8'5
ζ Persei (1)	Nov.—Mar.	3 47'0	+31 32	12'8, 32'6	208, 287	3'0, 8'7, 11'1
9 (Hevel) Camelopardi	Oct.—Apr.	3 47'4	+60 46	2'2	41	5'2, 8'2
30 Eridani	Dec.—Feb.	3 47'1	-5 42	8'2	135	6'5, 10'5
Pair in Perseus	Nov.—Mar.	3 49'5	+41 33	8'8	149	6'7, 9'7
Pair in Cepheus (2)	—	3 50'9	+80 23	0'96	35	5'7, 7'0
Pair in Perseus	Nov.—Mar.	3 52'1	+38 29	1'60	329	6'2, 9'7
Pair in Perseus (3)	Nov.—Mar.	3 59'7	+33 9	1'08	204	6'7, 9'0
P. iii. 242 Persei (4)	Nov.—Mar.	4 0'0	+37 47	2'7	138	6'8, 9'5
P. iv. 53 Tauri	Nov.—Mar.	4 15'7	+20 30	2'1	171	5'6, 8'8
56 Persei	Nov.—Mar.	4 17'2	+33 42	4'4	50	5'5, 9'2
2 Camelopardi	Nov.—Apr.	4 30'9	+53 15	1'62	292	5'8, 7'5
7 Camelopardi (5)	Nov.—Apr.	4 48'2	+53 34	1'24, 25'7	309, 239	4'6, 7'9, 11'0
5 Aurigæ	Oct.—Apr.	4 52'5	+39 13	2'7	247	6'0, 9'7
14 Orionis	Dec.—Mar.	5 1'6	+8 20	1'15	203	5'8, 6'0
16 Aurigæ (6)	Nov.—Apr.	5 10'7	+33 15	4'3	57	5'0, 10'6
Leporis 28	Dec.—Feb.	5 15'7	-21 22	4'3	283	6'0, 10'5
Pair in Orion	Dec.—Mar.	5 18'6	-0 59	1'43	171	6'5, 6'7
η Orionis	Dec.—Mar.	5 18'7	-2 30	1'00	85	4'0, 6'0
ψ <sup>2</sup> Orionis	Dec.—Mar.	5 20'8	+2 59	2'7	324	5'4, 9'0
β Leporis	Dec.—Feb.	5 23'4	-20 51	2'5	285	3'5, 11'0
31 Orionis	Dec.—Mar.	5 23'9	-1 11	12'6	88	5'4, 10'5
33 Orionis	Dec.—Mar.	5 25'2	+3 12	1'76	28	6'0, 7'0
Pair in Taurus	Nov.—Mar.	5 30'1	+26 52	1'06	178	6'5, 7'0
42 Orionis	Dec.—Mar.	5 29'8	-4 55	1'73	218	5'2, 8'9
Pair in Camelopardus	Nov.—Apr.	5 37'5	+62 46	1'54	23	6'4, 7'3
Pair in Orion	Nov.—Mar.	5 41'5	+20 50	0'85, 75'6	315, 161	6'2, 8'0, 7'5
52 Orionis	Dec.—Mar.	5 41'9	+6 25	1'61	204	6'1, 6'5
Leporis 61	Dec.—Feb.	5 44'4	-14 31	2'7	179	6'0, 9'4
θ Aurigæ (7)	Nov.—Mar.	5 52'0	+37 12	2'4	359	3'0, 8'6
3 Monocerotis	Dec.—Mar.	5 56'5	-10 36	1'62	355	6'0, 9'7
Pair in Auriga	Nov.—Mar.	5 59'7	+36 17	1'73	277	7'0, 10'0
4 Monocerotis	Dec.—Mar.	6 3'1	-11 7	3'2, 9'0	178, 244	6'7, 10'5, 11'5
Pair in Monoceros	Dec.—Mar.	6 6'1	-4 38	0'98	170	6'2, 8'7
4 Lyncis	Nov.—May	6 11'9	+59 25	0'95	101	6'2, 7'5
Pair in Camelopardus	—	6 15'3	+70 36	5'5	79	6'0, 10'9
Pair in Monoceros	Jan.—Apr.	6 16'1	-11 42	3'8	26	6'2, 9'9
54 Aurigæ	Jan.—Apr.	6 32'4	+28 22	0'83	37	6'0, 8'0
15 Monocerotis (8)	Jan.—Apr.	6 34'7	+10 0	3'0, 16'9	211, 13	Var. 8'7, 11'0
Canis Maj. 29	Feb.—Apr.	6 43'8	-15 1	0'97	290	6'0, 8'7

(1) A 9'3 at 108°, 89" 1, and a 10'0 at 185°: 119" 5.

(2) Binary.

(3) A 12'5 mag. at 119°: 34" 7.

(4) Rapid common proper motion, in which 50 Persei, 12' distant, joins. Parallax insensible.

(5) 8'7 mag. of a very dusky hue.

(6) A pair with double companion in field, 32° J. and 10' north of 16.

(7) 8'6 possibly variable. Several faint distant companions.

(8) In the midst of a scattered cluster, containing several pairs.



NAME OF OBJECT	Visible	R.A.	Decl.	Distances	Posit. Angles	Magnitude
15 <b>Lyncis</b> (1) . . . .	Nov.—May	<sup>h. m.</sup> 6 47'4	+58° 35'	0"73	360°	5'0, 7'3
λ <b>Geminorum</b> . . . .	Dec.—Apr.	7 11'5	+16 44	9'5	33	3'5, 9'8
P. vii. 52 <b>Canis Minor</b> . . . .	Dec.—Apr.	7 11'5	+ 9 30	1'28	113	7'0, 7'0
30 <b>Canis Maj.</b> . . . .	Jan.—Mar.	7 14'0	-24 45	7'8, 14'3	90, 80	6'0, 10'5, 11'2
P. vii. 116 <b>Monocerotis</b> . . . .	Jan.—Apr.	7 22'5	-11 20	0'80, 2'0, 23'4	166, 313, 157	6'0, 8'2, 8'9, 10'0
Pair in <b>Gemini</b> . . . .	Dec.—May	7 27'9	+31 12	0'82	332	5'5, 6'5
π <b>Geminorum</b> . . . .	Dec.—May	7 40'1	+33 42	22'0	212	5'0, 10'8
Pair in <b>Canis Minor</b> . . . .	Dec.—May	7 46'7	+ 3 41	1'20	43	7'0, 7'2
Pair in <b>Cancer</b> . . . .	Dec.—May	7 54'2	+23 54	2'7	333	6'1, 10'7
11 <b>Cancri</b> . . . .	Dec.—May	8 1'8	+27 49	3'2	218	6'9, 10'4
Pair in <b>Lynx</b> . . . .	Dec.—June	8 7'0	+43 23	4'2	294	6'7, 10'5
Pair in <b>Argo</b> (2) . . . .	Feb.—May	8 34'0	-19 20	4'3	104	6'5, 10'5
<b>Pyxidis</b> 17 . . . .	Feb.—May	8 34'2	-22 17	1'37	34	6'0, 9'0
α <b>Cancri</b> . . . .	Dec.—May	8 47'3	+31 1	1'42	328	5'8, 6'2
ι <b>Ursæ Maj.</b> (3) . . . .	Dec.—June	8 51'4	+48 29	9'6	357	3'2, 10'2
σ² <b>Ursæ Maj.</b> (4) . . . .	Dec.—June	9 0'4	+67 36	2'4	233	5'1, 8'7
37 <b>Lyncis</b> . . . .	Dec.—June	9 12'8	+51 44	5'5	117	6'0, 10'6
κ <b>Leonis</b> . . . .	Dec.—May	9 18'0	+26 41	3'4	205	5'0, 10'2
ω <b>Leonis</b> (5) . . . .	Jan.—June	9 22'4	+ 9 33	0'68	100	6'0, 6'4
3 <b>Leonis</b> . . . .	Jan.—June	9 22'4	+ 8 41	25'1	79	6'0, 10'8
Pair in <b>Leo</b> . . . .	Jan.—June	9 32'6	+11 17	8'3	103	6'7, 10'2
<b>Felis</b> 15 . . . .	Mar.—June	9 36'2	-17 58	3'1	261	7'2, 11'0
P. x. 23 <b>Leonis</b> . . . .	Feb.—June	10 10'0	+18 18	0'76	217	6'5, 7'5
<b>Felis</b> 54 . . . .	Mar.—June	10 16'2	-21 58	2'0	189	6'5, 9'0
P. x. 94 <b>Sextantis</b> . . . .	Mar.—June	10 25'3	- 7 3	2'8	166	6'0, 9'9
Pair in <b>Ursa Maj.</b> . . . .	Feb.—June	10 41'5	+41 42	0'84	330	6'5, 7'5
ψ <b>Crateris</b> . . . .	Apr.—June	11 6'9	-17 53	0'45	149	6'2, 6'9
ν <b>Ursæ Maj.</b> . . . .	Feb.—June	11 12'3	+33 43	7'0	147	3'8, 9'6
γ <b>Crateris</b> (6) . . . .	Apr.—June	11 19'2	-17 4	5'1	98	4'0, 10'0
Pair in <b>Ursa Maj.</b> (7) . . . .	Apr.—Aug.	11 25'8	+61 43	1'26	65	6'3, 7'7
P. xi. 126 <b>Virginis</b> . . . .	Mar.—June	11 32'6	- 1 48	4'9	279	6'2, 9'7
Pair in <b>Can. Ven.</b> . . . .	Mar.—June	11 50'4	+36 5	1'55	122	6'5, 9'0
Pair in <b>Can. Ven.</b> . . . .	Mar.—June	12 5'0	+40 32	1'21	338	6'2, 7'0
Pair in <b>Comæ Ber.</b> . . . .	Mar.—June	12 11'8	+29 34	8'2	277	6'2, 10'5
P. xii. 104 <b>Corvi</b> . . . .	Apr.—June	12 24'2	-12 45	1'81	354	6'2, 10'2
Pair in <b>Corvus</b> . . . .	Apr.—June	12 29'8	-16 12	11'2	258	6'7, 11'2
35 <b>Comæ Ber.</b> (8) . . . .	Apr.—July	12 47'7	+21 52	1'37, 28'8	68, 125	5'2, 8'0, 9'2
46 <b>Virginis</b> . . . .	Apr.—June	12 54'7	- 2 45	1'44	151	5'4, 9'5
<b>Hydræ</b> 348 . . . .	Apr.—June	12 57'6	-19 58	0'71	133	6'2, 6'2
P. xii. 268 <b>Can. Ven.</b> . . . .	Apr.—June	13 0'7	+29 38	6'5, 40'3	219, 7	6'0, 10'5, 12'5

(1) Binary.  
 (2) Red, blue.  
 (3) Very large common proper motion, in which a neighbouring star,  $\iota$  Ursæ, participates.  
 (4) Binary, closing rapidly.

(5) Binary, period about 115 years.  
 (6) Common proper motion.  
 (7) Binary, period 94½ years.  
 (8) Binary.

NAME OF OBJECT	Visible	R.A.	Decl.	Distances	Posit. Angles	Magnitude
Pair in Can. Ven. (1)	Apr.—June	h. m. 13 6'7	+32° 41'	1'41"	348°	6'3, 6'9
25 Can. Ven. (2)	Apr.—June	13 32'4	+36 52	0'80	151	6'2, 8'2
7 Boötis (3)	Apr.—June	13 41'8	+18 1	8'9	353	4'1, 11'5
P. xiii. 242 Can. Ven.	Apr.—June	13 49'3	+30 28	1'81	18	7'0, 9'6
Pair in Boötes	Apr.—June	14 1'8	+35 19	14'2	70	6'4, 10'0
P. xiv. 20 Boötis	May—July	14 8'3	+12 31	1'95	249	6'6, 9'3
♄ Virginis	May—July	14 22'3	- 1 43	4'1	112	5'2, 9'4
Pair in Boötes	May—Aug.	14 28'5	+49 41	5'0	131	7'2, 11'0
Pair in Boötes	May—Aug.	14 36'3	+49 11	7'0	102	7'3, 11'0
Libræ 23	May—June	14 42'1	-16 52	1'27	236	7'0, 8'0
Pair in Boötes	May—July	14 43'4	+24 50	1'50	55	6'3, 7'4
Pair in Boötes	May—July	14 48'0	+16 10	1'49	195	6'2, 7'2
Pair in Boötes	May—July	14 51'3	+32 45	4'3	113	6'2, 10'5
Pair in Boötes	May—Aug.	15 10'6	+38 44	1'38	257	6'2, 8'2
Pair in Libra (4)	May—July	15 12'5	-23 51	1'90	180	7'0, 9'0
5 Serpenteis (5)	May—July	15 13'4	+ 2 13	10'7	38	4'8, 10'0
Coronæ 1 (6)	May—July	15 13'5	+27 15	1'22	308	5'6, 6'1
6 Serpenteis	May—July	15 15'2	+ 1 8	2'3	13	4'7, 9'4
♁ Coronæ Bor. (7)	May—Aug.	15 18'5	+30 42	0'65	182	5'7, 6'0
μ <sup>a</sup> Boötis (8)	May—Aug.	15 20'3	+37 45	0'78	104	6'5, 7'8
2 Scorpionis	May—June	15 46'7	-25 0	2'6	279	5'5, 9'0
11 Scorpionis	May—July	16 1'3	-12 26	3'3	256	6'1, 10'4
Pair in Cor. Bor.	May—Aug.	16 7'3	+33 38	5'4	262	6'0, 10'5
Pair in Cor. Bor.	May—July	16 8'0	+26 58	2'8	136	6'2, 10'7
Pair in Serpens	May—July	16 16'0	+ 1 26	2'6	1	7'2, 9'9
η Draconis	Apr.—Aug.	16 22'4	+61 46	5'2	142	2'8, 9'0
Draconis 99	Apr.—Aug.	16 22'3	+61 57	1'16	2	6'2, 7'4
Pair in Hercules	May—Aug.	16 23'3	+26 15	1'26	211	6'6, 7'9
ζ Herculis (9)	May—Aug.	16 37'0	+31 49	1'65	82	2'6, 7'0
Pair in Hercules	May—July	16 44'3	+13 27	5'4	39	5'7, 10'3
21 Ophiuchi	June—Aug.	16 45'6	+ 1 25	0'98	162	6'0, 8'0
52 Herculis	May—Aug.	16 46'0	+46 11	1'83	309	5'0, 10'0
P. xvi. 270 Oph.	June—Aug.	16 56'5	+ 8 36	1'28	157	6'7, 7'9
Herculis 206 (10)	May—Aug.	16 59'8	+19 45	1'78	232	6'9, 10'2
P. xvii. 18 Oph.	June—Aug.	17 7'5	+ 7 54	1'47	204	6'7, 8'9
P. xvii. 43 Oph.	July—Aug.	17 13'2	-17 38	1'74	261	6'2, 7'5
68 Herculis	June—Aug.	17 13'3	+33 14	4'4	61	5'1, 10'1
Pair in Hercules	June—Sept.	17 42'1	+17 44	0'79	298	6'1, 7'8
Pair in Taurus Pon.	June—Sept.	17 42'2	+39 22	7'6	350	6'7, 10'3
90 Herculis	June—Sept.	17 49'6	+40 3	1'90	122	5'9, 9'4

(1) Binary.

(2) Binary, period 120 years.

(3) Large common proper motion.

(4) A pair in field, 31<sup>a</sup> f., 7<sup>b</sup>, 7<sup>c</sup> : 329° : 0''80.

(5) Large common proper motion.

(6) Magnitudes possibly variable.

(7) Binary, period 41½ years.

(8) Binary, period rather uncertain.

(9) Binary, period 34½ years.

(10) Pair in field, 47<sup>a</sup> f., 5<sup>a</sup> n. 6'9, 11'3 : 228° : 1''50.

NAME OF OBJECT	Visible	R.A.	Decl.	Distances	Posit. Angles	Magnitude
Pair in Hercules	June—Oct.	h. m.				
Herculis 417	June—Sept.	18 4'2	+49 42	2'2	148°	6'4, 10'5
μ Sagittarii (1)	June—Aug.	18 5'1	+16 27	1'19	236	6'7, 7'8
Pair in Scutum Sob.	June—Aug.	18 6'9	-21 5	16'8	258	3'5, 11'0
21 Sagittarii	June—Aug.	18 15'7	-15 9	12'4, 1'27	220, 64	7'0, 8'2, 8'5
	June—Aug.	18 18'6	-20 36	2'1	293	5'2, 8'0
Pair in Lyra	June—Nov.	18 32'4	+33 22	7'3	205	5'6, 10'5
Pair in Aquila	July—Oct.	18 33'0	+ 4 15	1'23	289	6'6, 9'5
Lyrae 91	June—Nov.	18 50'7	+33 49	1'86, 45'3	134, 350	6'0, 10'0, 7'3
17 Lyrae	June—Nov.	19 3'1	+32 20	3'7	321	5'5, 9'7
Pair in Lyra	June—Nov.	19 11'3	+27 16	0'83	156	6'6, 7'2
Pair in Cygnus	June—Nov.	19 12'3	+49 52	2'3	76	6'9, 10'4
2 Vulpeculae	July—Nov.	19 12'9	+22 49	1'86	125	5'7, 9'5
P. xix. 108 Drac.	Feb.—Nov.	19 15'7	+63 0	1'16	337	7'0, 8'1
Pair in Cygnus	July—Dec.	19 39'0	+40 27	0'85	25	6'5, 7'9
P. xix. 263 Cygni	July—Dec.	19 39'7	+38 3	0'96	196	7'0, 8'0
δ Cygni (2)	July—Dec.	19 41'4	+44 50	1'68	318	2'8, 7'5
π Aquilae	July—Oct.	19 43'4	+11 32	1'43, 31'2	117, 306	6'1, 6'7, 11'0
16 Vulpeculae	July—Nov.	19 57'2	+24 37	0'69	95	5'7, 5'9
Cygni 153	July—Dec.	20 9'3	+51 7	4'0	81	5'9, 10'9
Pair in Cygnus	July—Dec.	20 10'4	+41 45	0'90, 11'8	172, 34	7'0, 7'6, 9'3
π Capricorni	July—Sept.	20 20'8	-18 35	3'3	145	5'1, 8'7
Pair in Vulpecula	Aug.—Nov.	20 27'1	+25 24	1'16	78	6'3, 7'6
Delphini 43	Aug.—Nov.	20 39'5	+11 54	1'22, 38'7	92, 34	6'4, 8'0, 12'0
13 Delphini	Sept.—Nov.	20 42'2	+ 5 35	1'61	186	5'2, 8'9
λ Cygni (3)	Aug.—Dec.	20 43'0	+36 4	0'65, 85'0	80, 105	5'0, 7'0, 8'7
60 Cygni	Aug.—Dec.	20 57'2	+45 42	2'7	165	5'5, 9'5
γ Equulei (4)	Aug.—Nov.	21 4'9	+ 9 38	2'2, 41'3	275, 10	4'3, 10'0, 12'0
P. xxi. 51 Cephei	June—Jan.	21 9'0	+59 31	1'12	226	5'9, 6'6
P. xxi. 50 Cygni	Aug.—Dec.	21 9'9	+40 41	1'40	128	6'6, 7'1
7 Cygni (5)	Aug.—Dec.	21 10'2	+37 33	1'10	116	3'7, 7'8
Pair in Cepheus	June—Jan.	21 11'6	+63 59	0'95	248	6'6, 6'9
P. xxi. 166 Cephei	June—Jan.	21 24'3	+59 16	12'2	190	6'2, 11'0
Pair in Cepheus	June—Jan.	21 52'6	+64 47	1'45	246	6'0, 8'5
Pair in Pegasus	Sept.—Dec.	21 54'6	+23 24	2'9	259	6'6, 11'1
15 Cephei	June—Feb.	22 0'6	+59 16	10'9	297	6'5, 11'0
Pair in Pegasus	Sept.—Dec.	22 8'8	+ 7 25	1'20	126	6'0, 7'7
Pair in Cepheus	—	22 29'8	+69 20	0'84	265	6'5, 7'0
P. xxii. 258 Cephei	—	22 48'0	+82 33	3'6	34	5'1, 10'3
Pair in Lacerta	Sept.—Dec.	22 48'6	+44 9	1'33, 26'4	217, 354	6'0, 8'0, 10'7
Pair in Pegasus	Sept.—Dec.	22 51'2	+11 14	3'6	10	6'4, 9'1

(1) A 9'5 at 312° : 48''3, and another at 115° : 50''1, a 12'5 mag. at 118° : 25''2, and an excessively faint star at 180° ± 25''; discovered by Mr. Common with 37-in. reflector. Good test for large apertures.

(2) Binary, distance constant.  
 (3) Binary.  
 (4) 4'3 and 10'0 moving together through space.  
 (5) Binary.

NAME OF OBJECT	Visible	R.A.	Decl.	Distances	Posit. Angles	Magnitude
52 Pegasi . . . . .	Sept.—Dec.	h. m. 22 53'5	+11 7	" 1'22	213°	6'0, 8'0
π Cephei (1) . . . . .	—	23 4'3	+74 46	1'31	27	4'7, 8'7
Pair in Cassiopeia . . . . .	Aug.—Mar.	23 5'2	+56 50	1'54	305	6'9, 9'0
96 Aquarii (2) . . . . .	Oct.—Dec.	23 13'5	— 5 45	9'9	23	6'2, 11'3
P. xxiii. 101 Cass. (3) . . . . .	Aug.—Mar.	23 24'7	+57 55	1'38	345	5'0, 9'4
ω <sup>3</sup> Aquarii . . . . .	Oct.—Dec.	23 36'8	—15 10	5'7	88	5'0, 11'0
78 Pegasi . . . . .	Sept.—Jan.	23 38'3	+28 44	1'46	192	5'0, 8'1
Pair in Pegasus . . . . .	Oct.—Jan.	23 39'7	+19 47	1'98	62	6'7, 9'3
Pair in Andromeda . . . . .	Sept.—Jan.	23 52'5	+34 23	3'4	23	6'4, 9'1
Pair in Andromeda . . . . .	Sept.—Jan.	23 58'7	+41 27	5'2	168	6'1, 10'0

(1) Binary.  
(2) Common proper motion.  
(3) An 8'0 mag. at 269°:75''9 is double, 7'5, 9'5: 221°:1''36, with a companion, 11'6 mag. at 337°:26''9. The 5'0 has a 10'1 mag. companion at 115°:43''5, and another at 339°:66''9 which is itself double, having a 10 mag. star at 74°:10''9.

## SHOOTING STARS.

## Radiant Points.

DATE	Radiant Point		NOTES	Radiant Point	
	R.A.	Decl.		Rises	Souths
January 2-3 . . . . .	h. m. XV. 30	+48°	Fine morning shower.*	—	8 39 A.M.
March 7 . . . . .	XVI. 16	+15	Formerly active.	9 51 P.M.	5 13 A.M.
April 19-20 . . . . .	XVIII. 1	+33	<i>Lyrids</i> .	6 27 P.M.	4 8 A.M.
April 29-May 2 . . . . .	XXI. 45	— 2	Morning shower. Max., May 2.	1 15 A.M.	7 8 A.M.
July 30-August 1 . . . . .	II. 8	+53	Fine display, 1878.*	—	5 29 A.M.
August 9-10 . . . . .	II. 52	+56	<i>Perseids</i> . Very fine annual shower.	—	5 38 A.M.
September 1 . . . . .	XX. 20	+54	Large meteors.* [10th max.*	—	9 36 P.M.
October 17-20 . . . . .	VI. 0	+15	Fine annual shower.	8 47 P.M.	4 10 A.M.
November 13-14 . . . . .	X. 0	+23	Magnificent shower in 1866. <i>Leonid</i> .	10 14 P.M.	6 27 A.M.
November 19-23 . . . . .	IV. 16	+20	Long-continued shower.	4 19 P.M.	0 12 A.M.
November 27 . . . . .	I. 40	+43	<i>Andromedes</i> . Fine display 1872-85.*	—	9 13 P.M.
December 9-12 . . . . .	VII. 0	+32	<i>Geminids</i> . Rich shower.	4 10 P.M.	1 41 A.M.

\* The radiant point of these showers is circumpolar, and never sets.



## TEST OBJECTS.

Two Inches Aperture.

NAME OF OBJECT	Visible	R.A.	Decl.	Distances	Posit. Angles	Magnitude
DIVIDING TESTS						
		h. m.	o ' "	"	o	
$\epsilon^1$ Lyræ . . . . .	June—Nov.	18 40'6	+39 33	3'12	16'1	4'6, 6'3
$\Sigma$ 2671 . . . . .	May—Dec.	20 15'6	+55 2	2'89	341'2	6'0, 7'2
$\Sigma$ 389 . . . . .	Sept.—Mar.	3 20'9	+58 58	2'71	66'5	6'4, 7'6
P. xvii. 300 . . . . .	June—Sept.	17 51'5	+18 21	2'65	115'1	6'9, 7'0
$\gamma$ Arietis . . . . .	Oct.—Mar.	1 43'7	+21 42	2'65	168'5	6'1, 7'0
P. xiii. 238 . . . . .	April—June	13 49'0	- 7 30	2'64	75'0	6'7, 7'3
$\delta$ 1 Virginis . . . . .	April—June	13 31'6	- 7 18	2'63	42'3	7'0, 7'3
$\Sigma$ 425 . . . . .	Oct.—Feb.	3 32'9	+33 45	2'60	97'1	7'0, 7'1
$\epsilon^2$ Lyræ . . . . .	June—Nov.	18 40'6	+39 30	2'58	136'7	5'1, 5'2
$\mu$ Draconis . . . . .	May—Oct.	17 3'0	+54 37	2'40	163'0	5'0, 5'0
$\Sigma$ 899 . . . . .	Nov.—Feb.	3 16'1	+17 38	2'34	20'2	7'0, 7'5
$\Sigma$ 2950 . . . . .	July—Jan.	22 46'9	+61 6	2'27	311'7	5'7, 6'9
DEFINING TESTS						
$\eta$ Cassiop. . . . .	Aug.—Feb.	0 42'1	+57 13	4'90	175'0	3'5, 7'3
$\delta$ Argûs . . . . .	Dec.—April	7 42'6	-11 55	3'28	17'1	5'7, 7'5
$\delta$ Serpentis . . . . .	May—July	15 29'4	+10 55	3'48	188'4	3'9, 5'5
$\delta$ 2 Aquarii . . . . .	July—Nov.	20 58'1	- 6 16	2'76	191'6	6'0, 8'0
$\epsilon$ Boötis . . . . .	May—July	14 40'0	+27 33	3'02	329'5	2'5, 5'0
$\gamma$ Ceti . . . . .	Nov.—Jan.	2 37'4	+ 2 45	2'93	289'7	3'4, 7'1
SPACE-PENETRATING TESTS						
$\alpha$ Ursæ Min. ( <i>Polaris</i> ) . . . . .	—	1 17'0	+88 42	18'3	213'8	2'0, 8'4
$\beta$ Orionis ( <i>Rigel</i> ) . . . . .	Dec.—Mar.	5 9'2	- 8 20	9'5	201'1	1'0, 7'8
$\gamma^1$ Aquarii . . . . .	Aug.—Nov.	22 40'7	-14 40	28'5	114'6	5'9, 9'2
$\delta$ 1 Aquilæ . . . . .	June—Sept.	18 53'9	+13 28	17'0	260'0	5'1, 9'0
$\delta$ 2 Ophiuchi . . . . .	June—Sept.	16 41'4	+ 2 16	22'3	91'9	6'0, 9'0
P. xii. 221 . . . . .	April—June	12 49'8	+12 7	28'6	203'0	6'8, 9'0
Camelop. 4 . . . . .	Aug.—Feb.	3 21'3	+55 3	14'6	159'2	5'4, 9'4
Tauri 39 . . . . .	Nov.—Mar.	3 34'4	+ 4 46	26'2, 37'7	55'7, 301'2	5'6, 9'0, 9'4
Camelop. 176 . . . . .	—	8 8'1	+72 46	43'9	85'5	5'7, 9'3
$\delta$ 2 Ursæ Maj. . . . .	Dec.—June	9 22'5	+63 33	22'8	271'5	4'0, 9'2
Scut. Sob. (29) . . . . .	July—Sept.	18 25'1	-10 53	12'2	256'5	6'0, 9'2
$\alpha$ Cassiop. . . . .	Aug.—Feb.	0 34'0	+55 55	62'4	279'8	2'4, 9'5

## Two and a Half Inches Aperture.

NAME OF OBJECT	Visible	R.A.	Decl.	Distances	Posit. Angles	Magnitude
DIVIDING TESTS						
		h. m.	o.	"	°	
Tauri. Pon. 75 . . . . .	June—Sept.	18 39'9	+ 5 22	2'22	115'3	6'2, 6'6
P. xx. 429. . . . .	June—Dec.	20 54'8	+50 1	2'06	31'9	6'2, 7'0
P. o. 181 . . . . .	Aug.—Feb.	0 41'5	+50 50	2'05	147'4	7'0, 7'8
Cephei 287 . . . . .	—	23 23'1	+73 29	2'03	32'6	7'0, 8'0
85 Lynceis . . . . .	Jan.—Apr.	8 2'3	+32 30	2'01	47'6	7'1, 7'9
Ursæ Maj. 234 . . . . .	Apr.—Aug.	11 32'4	+64 59	1'97	322'7	6'7, 7'7
84 Aurigæ . . . . .	Dec.—May	6 30'6	+41 41	1'94	81'1	6'9, 7'7
Σ 2624 . . . . .	June—Nov.	19 59'2	+35 42	1'92, 42'3	175'5, 327'5	7'0, 7'6, 9'5
02 358 . . . . .	June—Sept.	18 30'8	+16 53	1'88	19'4	6'5, 6'8
γ Ophiuchi . . . . .	June—Aug.	17 56'9	— 8 11	1'84	254'5	5'0, 6'0
Σ 1116 . . . . .	Dec.—Apr.	7 28'1	+12 33	1'82	109'9	7'0, 7'7
Σ 1871 . . . . .	Apr.—July	14 37'7	+51 53	1'80	291'5	7'0, 7'3
DEFINING TESTS						
μ Can. Maj. . . . .	Dec.—Mar.	6 50'9	—13 54	2'92	338'8	5'2, 8'2
38 Lynceis . . . . .	Feb.—June	9 11'8	+37 18	2'80	239'1	4'2, 6'3
* Draconis . . . . .	—	19 48'6	+69 59	2'94	5'5	4'2, 7'2
* Leporis . . . . .	Dec.—Mar.	5 8'0	—13 5	2'41	357'4	5'2, 7'7
* Hydræ . . . . .	Jan.—Apr.	8 40'7	+ 6 50	3'34	228'0	3'8, 7'4
* Leonis . . . . .	Mar.—June	11 17'9	+11 9	2'80	63'0	4'6, 7'4
SPACE-PENETRATING TESTS						
o Persei . . . . .	Nov.—Mar.	3 37'1	+31 55	20'0	238'4	4'5, 9'0
66 Eridani . . . . .	Dec.—Mar.	5 1'1	— 4 49	52'5	9'4	6'0, 9'2
β Serpentis . . . . .	May—July	15 41'0	+15 47	30'7	265'0	3'6, 9'1
33 Arietis . . . . .	Oct.—Mar.	2 34'0	+26 35	28'6	359'6	5'2, 9'3
α Lyræ (Vega) . . . . .	June—Nov.	18 33'1	+38 41	49'2	155'5	0'2, 9'5
12 Lacertæ . . . . .	July—Nov.	22 36'4	+39 38	70'5	15'9	5'7, 9'7
P. v. 37 . . . . .	Nov.—Mar.	5 12'4	+20 0	9'0	203'0	6'0, 9'8
Cassiop. 63 . . . . .	Aug.—Feb.	0 32'4	+46 20	10'4	85'9	6'8, 9'9
θ Virginis . . . . .	Apr.—June	13 4'0	— 4 56	7'1, 70'9	344'6, 297'1	4'9, 8'5, 10'0
18 Libræ . . . . .	May—July	14 52'7	—10 41	19'5	38'9	6'0, 10'0
P. xix. 144 . . . . .	July—Oct.	19 24'5	+ 2 39	34'2	5'2	6'0, 10'0
μ Herculis . . . . .	May—Sept.	17 42'0	+27 47	31'2	243'9	3'5, 10'0

Three Inches Aperture.

NAME OF OBJECT	Visible	R.A.	Decl.	Distances	Posit. Angles	Magnitude
DIVIDING TESTS						
		h. m.	o ' "	"	o	
33 Orionis . . . . .	Dec.—Mar.	5 25'2	+ 3 12	1'76	28'3	6'0, 7'0
12 Lyncis . . . . .	Dec.—May	6 36'2	+59 33	1'70, 8'7	126'8, 307'5	5'7, 6'4, 7'4
Σ 1333 . . . . .	Feb.—June	9 11'4	+35 51	1'69	42'2	6'7, 7'0
Persei 85 . . . . .	Aug.—Feb.	2 44'8	+52 31	1'69	301'8	6'8, 7'1
OS 437 . . . . .	July—Nov.	21 16'1	+31 58	1'64	49'3	6'0, 6'5
λ Ophiuchi . . . . .	June—Aug.	16 25'2	+ 2 14	1'65	44'5	4'4, 5'4
Cephei 83 . . . . .	May—Dec.	20 59'0	+56 13	1'62	349'3	6'4, 6'8
Lyncis 157 . . . . .	Feb.—June	9 13'9	+38 41	1'59	154'8	6'8, 7'4
Σ 644 . . . . .	Oct.—Mar.	5 2'6	+37 9	1'58	220'9	6'8, 7'1
Σ 2744 . . . . .	July—Oct.	20 57'3	+ 1 5	1'54	170'0	6'3, 7'0
μ Libræ . . . . .	May—July	14 43'1	-13 41	1'51	337'4	5'2, 6'2
Ceti 187 . . . . .	Nov.—Feb.	1 14'3	-16 24	1'50	20'0	7'1, 7'3
DEFINING TESTS						
o Cephei . . . . .	July—Mar.	23 13'9	+67 29	2'59	192'6	5'2, 7'6
49 Leonis . . . . .	Feb.—June	10 29'0	+ 9 14	2'39	157'5	6'2, 8'4
P. xx. 376 . . . . .	July—Oct.	20 50'0	+ 4 6	1'97	287'5	6'2, 7'9
70 Ophiuchi . . . . .	June—Aug.	17 59'7	+ 2 32	2'05	20'5	4'3, 6'2
84 Virginis . . . . .	Apr.—June	13 37'3	+ 4 7	3'56	234'3	5'7, 8'0
23 Aquilæ . . . . .	July—Oct.	19 12'7	+ 0 52	3'33	11'6	5'7, 9'0
SPACE-PENETRATING TESTS						
59 Aurigæ . . . . .	Dec.—Apr.	6 45'2	+39 1	22'2	223'5	6'7, 10'0
λ Geminorum . . . . .	Dec.—Apr.	7 11'5	+16 45	9'5	33'0	3'5, 9'8
57 Pegasi . . . . .	Sept.—Jan.	23 3'7	+ 8 4	32'8	197'8	5'2, 10'0
P. xix. 307 . . . . .	July—Oct.	19 46'8	+10 4	13'3	309'7	6'6, 10'2
P. xix. 128 . . . . .	June—Nov.	19 21'5	+19 39	23'7, 68'7	40'9, 319'9	5'1, 10'1, 9'8
ν Ursæ Maj. . . . .	Feb.—June	11 12'3	+33 43	7'0	147'3	3'5, 9'6
δ Equulei . . . . .	July—Nov.	21 9'0	+ 9 33	40'0	21'2	4'5, 10'0
5 Serpentis . . . . .	May—July	15 13'4	+ 2 13	10'7	38 0	4'8, 10'0
31 Orionis . . . . .	Dec.—Mar.	5 23'9	- 1 11	12'6	87'8	5'4, 10'5
α Tauri (Aldebaran) . . . . .	Oct.—Feb.	4 29'4	+16 17	115'0	34'2	1'1, 10'3
θ Hydræ . . . . .	Feb.—May	9 8'4	+ 2 48	52'7	175'2	4'0, 10'5
β Delphini . . . . .	July—Nov.	20 32'2	+14 12	35'5	334'2	3'5, 10'6

## Three and a half Inches Aperture.

NAME OF OBJECT	Visible	R.A.	Decl.	Distances	Posit. Angles	Magnitude
DIVIDING TESTS						
		h. m.	° '	"	°	
Σ 1884 . . . . .	May—July	14 43'4	+24 50	1'50	55'0	6.3, 7.4
36 Androm. . . . .	Sept.—Feb.	0 48'9	+23 1	1'46	1'5	6.0, 6.4
42 Ceti . . . . .	Dec.—Feb.	1 14'0	- 1 6	1'45	355'0	6.7, 7.5
π Aquilæ . . . . .	July—Oct.	19 43'4	+11 32	1'43, 31'2	117'3, 306'0	6.1, 6.7, 11.0
♄ Cancri . . . . .	Dec.—May	8 47'3	+31 1	1'42	328'5	5.8, 6.2
0Σ 261 . . . . .	Apr.—July	13 6'7	+32 41	1'41	348'2	6.3, 6.9
P. xxi. 50 . . . . .	Aug.—Dec.	21 9'9	+40 41	1'40	127'7	6.6, 7.1
ε Arietis . . . . .	Dec.—Mar.	2 52'6	+20 54	1'39	201'3	5.4, 6.3
Pegasi 20 . . . . .	July—Oct.	21 23'3	+10 35	1'33	306'4	6.6, 6.6
P. vii. 170 . . . . .	Jan.—Mar.	7 34'1	+ 5 30	1'31	141'5	7.0, 7.3
Σ 1037 . . . . .	Jan.—Apr.	7 5'8	+27 25	1'30	310'6	6.9, 7.1
P. xvi. 270 . . . . .	June—Aug.	16 56'5	+ 8 36	1'29	157'0	6.7, 7.9
DEFINING TESTS						
6 Cassiop. . . . .	Aug.—Feb.	23 43'2	+61 35	1'60	195'6	5.1, 7.6
33 Pegasi . . . . .	July—Dec.	22 18'2	+20 17	2'06, 63'6	179'9, 328'6	6.1, 8.7, 8.3
ε Cassiop. . . . .	July—Mar.	2 19'7	+66 53	1'98, 7'4	262'0, 108'5	5.0, 7.5, 8.1
φ Virginis . . . . .	May—July	14 22'3	- 1 43	4'09	112'6	5.2, 9.4
P. xi. 126 . . . . .	Mar.—June	11 32'6	- 1 48	4'92	279'5	6.2, 9.7
17 Lyræ . . . . .	June—Nov.	19 3'1	+32 20	3'68	321'6	5.5, 9.7
SPACE-PENETRATING TESTS						
ε Leporis . . . . .	Dec.—Mar.	5 7'0	-12 0	12'6	336'2	4.5, 10.0
P. xx. 116 . . . . .	July—Oct.	20 18'8	+ 0 42	32'9	28'9	6.2, 10.3
θ Cancri . . . . .	Jan.—Apr.	8 25'0	+18 29	60'8	60'0	5.5, 10.4
υ Cygni . . . . .	June—Nov.	21 13'2	+34 25	15'1, 21'3	220'0, 178'5	4.3, 10.3, 10.2
α Androm. . . . .	Sept.—Feb.	0 2'5	+28 28	71'0	273'0	2.0, 10.4
13 Lacertæ . . . . .	July—Nov.	22 39'0	+41 13	14'7	129'4	5.1, 10.4
5 Lyncis . . . . .	Dec.—May	6 16'9	+58 29	30'3, 95'9	139'1, 272'5	6.0, 10.5, 8.5
γ Persei . . . . .	Aug.—Feb.	2 56'5	+53 3	57'7	323'7	3.1, 10.8
49 Piscium . . . . .	Sept.—Feb.	0 24'9	+15 24	16'7	105'8	7.0, 10.8
π Geminorum . . . . .	Jan.—Apr.	7 40'1	+33 42	22'0, 95'0	211'6, 340'0	5.0, 10.8, 10.5
56 Herculis . . . . .	May—Aug.	16 50'3	+25 55	18'0	93'1	6.0, 10.9
44 Virginis . . . . .	Apr.—June	12 53'8	- 3 12	20'9	54'6	5.8, 11.0



Four Inches Aperture.

Name of Object	Visible	R.A.	Decl.	Distances	Posit. Angles	Magnitude
DIVIDING TESTS						
		h. m.	o ' "	"	o	
P. vii. 52 . . . . .	Dec.—Apr.	7 11'5	+ 9 30	1'28	113'5	7'0, 7'0
‡ Scorp. . . . .	May—July	15 58'1	−11 4	1'27, 7'3	195'7, 66'0	5'0, 5'4, 7'2
Σ 2049 . . . . .	May—Aug.	16 23'3	+26 15	1'26	210'9	6'6, 7'9
Cor. Bor. 1 . . . . .	May—July	15 13'5	+27 15	1'22	308'1	5'6, 6'1
Σ 1606 . . . . .	Mar.—June	12 5'0	+40 32	1'21	338'1	6'2, 7'0
OΣ 182 . . . . .	Dec.—Apr.	7 46'7	+ 3 41	1'20	43'2	7'0, 7'2
Pegasi 148 . . . . .	Sept.—Dec.	22 8'8	+ 7 25	1'20	125'6	6'0, 7'7
Herculis 417 . . . . .	June—Sept.	18 5'1	+16 27	1'19	236'0	6'7, 7'8
Draconis 99 . . . . .	May—Oct.	16 22'3	+61 57	1'16	2'0	6'2, 7'4
14 Orionis . . . . .	Dec.—Mar.	5 1'6	+ 8 20	1'15	203'3	5'8, 6'0
Tauri Pon. 9 . . . . .	June—Aug.	17 51'2	+ 0 5	1'12	274'5	6'7, 7'0
P. xxi. 51 . . . . .	June—Jan.	21 9'0	+59 31	1'12	226'3	5'9, 6'6
DEFINING TESTS						
π Arietis . . . . .	Dec.—Mar.	2 42'9	+17 0	3'29, 25'2	122'5, 109'9	5'5, 8'2, 11'0
ζ Herculis . . . . .	May—Aug.	16 37'0	+31 49	1'65	82'0	2'6, 7'0
δ Cygni . . . . .	July—Dec.	19 41'4	+44 50	1'68	318'5	2'8, 7'5
ψ <sup>3</sup> Orionis . . . . .	Dec.—Mar.	5 20'8	+ 2 59	2'66	324'4	5'4, 9'0
5 Aurigæ . . . . .	Oct.—Apr.	4 52'4	+39 13	2'75	246'8	6'0, 9'7
η Draconis . . . . .	Apr.—Aug.	16 22'4	+61 46	5'26	142'1	2'8, 9'0
SPACE-PENETRATING TESTS						
ν <sup>1</sup> Cor. Bor. . . . .	May—Aug.	16 18'2	+33 58	66'4	236'6	5'1, 10'5
5 Ursæ Min. . . . .	—	14 27'8	+76 12	56'4	129'4	4'8, 10'5
3 Leonis . . . . .	Jan.—June	9 22'4	+ 8 41	25'1	79'2	6'0, 10'8
4 Ursæ Maj. . . . .	Jan.—June	8 51'4	+48 29	9'6	356'7	3'2, 10'2
41 Arietis . . . . .	Dec.—Mar.	2 43'4	+26 48	21'2, 34'0	265'8, 203'5	4'1, 11'3, 11'0
2 Lacertæ . . . . .	Aug.—Dec.	22 16'4	+45 58	48'2	9'7	5'0, 10'9
P. i. 145 . . . . .	Oct.—Feb.	1 34'9	+25 10	10'9	33'3	6'1, 10'9
42 Piscium . . . . .	Dec.—Feb.	0 16'5	+12 51	29'7	338'1	6'6, 11'0
54 Ophiuchi . . . . .	June—Aug.	17 29'1	+13 15	21'6	74'7	6'0, 11'0
κ Andromedæ . . . . .	Aug.—Feb.	23 34'8	+43 42	46'6, 103'2	188'7, 294'6	4'0, 11'0, 11'0
γ Libræ . . . . .	May—July	15 29'1	−14 25	41'3	151'8	4'5, 11'3
Lyncis 51 . . . . .	Dec.—Apr.	7 20'5	+48 26	16'9	94'3	6'2, 11'2

## Five Inches Aperture.

NAME OF OBJECT	Visible	R.A.	Decl.	Distances	Posit. Angles	Magnitude
DIVIDING TESTS						
		h. m.	o ' ,	"	o	
Vulpec. 94 . . . . .	Aug.—Nov.	20 27·1	+ 25 24	1·16	78·1	6·3, 7·6
Σ 115 . . . . .	Aug.—Feb.	1 16·1	+ 57 33	1·06	149·8	7·0, 7·0
Σ 749 . . . . .	Nov.—Mar.	5 30·1	+ 26 52	1·06	177·6	6·5, 7·0
η Orionis . . . . .	Dec.—Mar.	5 18·7	— 2 30	1·00	84·6	4·0, 6·0
OΣ 313 . . . . .	May—Sept.	16 28·7	+ 40 21	0·99	151·6	7·0, 7·6
21 Oph. . . . .	June—Aug.	16 45·6	+ 1 25	0·98	162·7	6·0, 8·0
OΣ 50 . . . . .	Aug.—Apr.	3 1·3	+ 71 7	0·98	213·4	7·0, 7·0
ε Equulei . . . . .	July—Nov.	20 53·4	+ 3 51	0·97, 10·5	285·6, 72·9	6·1, 6·8, 7·4
P. xix. 263 . . . . .	July—Dec.	19 39·8	+ 39 2	0·96	196·2	7·0, 8·0
48 Cephei . . . . .	—	3 50·9	+ 80 23	0·96	35·0	5·7, 7·0
4 Lynceis . . . . .	Nov.—May	6 11·9	+ 59 25	0·95	101·2	6·2, 7·5
A.C. 12 . . . . .	July—Oct.	19 52·4	— 2 34	0·93	332·4	7·0, 8·0
DEFINING TESTS						
ω <sup>a</sup> Aquarii . . . . .	Oct.—Dec.	23 36·8	— 15 10	5·68	87·8	5·0, 11·0
42 Orionis . . . . .	Dec.—Mar.	5 29·8	— 4 55	1·73	217·7	5·5, 9·2
13 Delphini . . . . .	Sept.—Nov.	20 42·2	+ 5 35	1·61	186·4	5·2, 8·9
60 Cygni . . . . .	Aug.—Dec.	20 57·2	+ 45 42	2·71	165·1	5·5, 9·5
68 Herculis . . . . .	June—Aug.	17 13·1	+ 33 14	4·41	61·8	5·1, 10·1
ψ Cassiop. . . . .	Aug.—Feb.	1 17·9	+ 67 32	28·7, 3·08	106·8, 256·0	4·5, 9·7, 10·6
SPACE-PENETRATING TESTS						
14 Aurigæ . . . . .	Nov.—Mar.	5 8·0	+ 32 33	11·9, 14·7	348·1, 225·4	5·1, 11·0, 7·5
14 Monoc. . . . .	Dec.—Apr.	6 28·6	+ 7 40	10·5	208·6	6·9, 10·9
40 Cassiop. . . . .	—	1 29·4	+ 72 27	53·3	237·0	6·0, 10·9
74 Oph. . . . .	June—Aug.	18 15·2	+ 3 19	27·9	285·6	5·0, 10·8
P. xxii. 36. . . . .	Aug.—Dec.	22 9·0	+ 39 9	27·2	178·5	6·0, 11·0
12 Ceti . . . . .	Dec.—Feb.	0 24·2	— 4 35	8·6, 212·4	187·0, 110·3	6·2, 10·9, 10·0
20 Pegasi . . . . .	Aug.—Nov.	21 55·5	+ 12 34	51·1	325·5	5·5, 11·4
55 Androm. . . . .	Aug.—Feb.	1 46·3	+ 40 10	60·1	355·2	5·5, 11·5
54 Sagittarii . . . . .	July—Sept.	19 34·2	— 16 32	35·8, 45·6	241·5, 41·7	5·5, 11·5, 9·0
96 Aquarii . . . . .	Oct.—Dec.	23 13·5	— 5 45	9·9	23·5	6·2, 11·3
85 Virginis . . . . .	Apr.—June	13 39·4	— 15 12	43·3	311·8	6·0, 11·7
56 Aquilæ . . . . .	Aug.—Oct.	19 47·9	— 8 52	46·7	77·8	5·7, 11·8

Six Inches Aperture.

NAME OF OBJECT	Visible	R.A.	Decl.	Distances	Posit. Angles	Magnitude
DIVIDING TESTS						
		h. m.	o. ' "	"	o	
Canis Maj. 89 . . . . .	Feb.—April	6 43'8	-15 1	0'97	288'9	6'0, 8'7
P. iv. 288 . . . . .	Oct.—Feb.	4 58'6	+19 38	0'88	336'8	6'5, 7'2
0Σ 175 . . . . .	Nov.—Mar.	5 27'9	+31 12	0'87	332'2	5'5, 6'5
Σ 2924 . . . . .	—	22 29'7	+69 20	0'84	265'6	6'5, 7'0
0Σ 229 . . . . .	Feb.—June	10 41'5	+41 42	0'84	330'1	6'5, 7'5
P. v. 222 . . . . .	Nov.—Mar.	5 41'5	+20 50	0'85, 75'6	314'7, 161'3	6'2, 8'0, 7'5
0Σ 383 . . . . .	July—Dec.	19 39'0	+40 27	0'85	25'1	6'5, 7'9
Σ 1883 . . . . .	April—July	14 43'2	+ 6 26	0'82	259'0	7'0, 7'0
0Σ 369 . . . . .	—	19 8'8	+71 54	0'80	41'3	7'0, 7'5
Σ 2215 . . . . .	June—Sept.	17 42'1	+17 44	0'79	300'3	6'1, 7'8
μ <sup>2</sup> Boötis . . . . .	May—Sept.	15 20'3	+37 47	108'4, 0'78	171'5, 104'0	4'5, 6'5, 7'8
β 132 . . . . .	July—Sept.	18 4'5	-19 52	0'79	237'0	6'7, 7'2
DEFINING TESTS						
Cygni 153 . . . . .	July—Dec.	20 9'3	+51 7	4'03	81'2	5'9, 10'9
θ Aurigæ . . . . .	Nov.—Mar.	5 52'0	+37 12	2'37	358'5	3'0, 8'6
κ Leonis . . . . .	Feb.—June	9 18'0	+26 41	3'36, 10	205'1, 65	5'1, 10'2, 11'5
π Cephei . . . . .	—	23 4'3	+74 46	1'31	27'3	4'7, 8'7
6 Serpentis . . . . .	May—July	15 15'2	+ 1 8	2'28	13'2	4'7, 9'4
Cancri 5 . . . . .	Dec.—April	7 54'2	+23 54	2'70	332'5	6'1, 10'7
SPACE-PENETRATING TESTS						
ε Ceti . . . . .	Dec.—Feb.	0 13'6	- 9 28	62'0	15'5	4'0, 11'5
ι Aquarii . . . . .	July—Oct.	20 33'6	+ 0 6	55'9, 72'9	217'4, 38'9	5'5, 11'5, 11'3
115 Tauri . . . . .	Dec.—April	5 20'5	+17 52	10'2	308'4	6'0, 11'4
Pegasi 129 . . . . .	Aug.—Nov.	22 4'8	+14 4	21'5	253'4	6'0, 11'8
72 Virginis . . . . .	April—June	13 24'5	- 5 53	29'2	16'1	6'2, 11'8
δ Cancri . . . . .	Jan.—April	8 38'2	+18 37	41'9	113'2	5'0, 11'8
41 Sextantis . . . . .	April—June	10 44'6	- 8 18	27'0	303'8	6'0, 11'8
ρ Boötis . . . . .	May—July	14 26'9	+30 52	53'3	334'0	3'6, 11'7
α <sup>2</sup> Capricorni . . . . .	Aug.—Oct.	20 11'7	-12 54	7'7	147'9	3'5, 11'5
ο Cassiop. . . . .	Aug.—Feb.	0 38'3	+47 40	32'2	303'9	5'0, 12'0
τ Aurigæ . . . . .	Nov.—April	5 41'3	+39 8	38'9, 47'9	350'0, 32'9	5'0, 12'0, 11'0
110 Herculis . . . . .	June—Sept.	18 40'8	+20 26	44'7, 61'2	95'6, 92'0	5'0, 12'0, 11'0

## Seven Inches Aperture.

NAME OF OBJECT	Visible	R. A.	Decl.	Distances	Posit. Angles	Magnitude
DIVIDING TESTS						
		h. m.	° ' "	"	°	
73 Ophiuchi . . . .	June—Aug.	18 3'9	+ 3 58	0'83	249'5	6'0, 7'6
54 Aurigæ . . . .	Jan.—Apr.	6 32'4	+28 22	0'83	36'9	6'0, 8'0
0Σ 338 . . . .	June—Sept.	17 46'8	+15 21	0'72	25'0	6'2, 6'4
Hydræ 348 . . . .	Apr.—June	12 57'6	-19 58	0'71	133'4	6'2, 6'2
Cygni 226 . . . .	July—Dec.	20 35'4	+40 10	0'70, 69'0	23'7, 69'8	6'4, 6'6, 7'6
16 Vulpeculæ . . . .	July—Nov.	19 57'2	+24 37	0'69	94'5	5'7, 5'9
Dawes 6 . . . .	Dec.—Mar.	5 23'3	- 3 24	0'69	84'4	7'1, 7'4
0Σ 156 . . . .	Dec.—Apr.	6 40'7	+18 19	0'68	317'1	6'6, 6'6
η Coronæ . . . .	May—Aug.	15 18'5	+30 42	0'65	182'2	5'7, 6'0
λ Cygni . . . .	Aug.—Dec.	20 43'0	+36 4	0'65	80'3	5'0, 7'0
B.A.C. 8277 . . . .	July—Mar.	23 43'2	+64 15	0'58, 48'9	253'8, 353'4	6'5, 7'7, 8'5
P. xviii. 132 . . . .	June—Sept.	18 30'8	+23 31	0'56	352'1	6'3, 6'6
DEFINING TESTS						
44 Cygni . . . .	July—Dec.	20 26'7	+36 33	2'26	157'9	6'3, 11'0
58 Ceti . . . .	Nov.—Feb.	1 52'2	- 2 37	2'73	12'8	6'3, 11'5
Lyræ 91 . . . .	June—Nov.	18 50'7	+33 49	1'86, 45'3	133'9, 350'6	6'0, 10'0, 7'3
τ Cygni . . . .	Aug.—Dec.	21 10'2	+37 33	1'10	116'3	3'7, 7'8
46 Virginis . . . .	Apr.—June	12 54'7	- 2 45	1'44	151'5	5'4, 9'5
46 Eridani . . . .	Dec.—Mar.	4 28'4	- 6 59	1'47	57'0	6'0, 10'0
SPACE-PENETRATING TESTS						
β Aquarii . . . .	Sept.—Nov.	21 25'6	- 6 4	34'3, 54'5	318'9, 184'9	3'0, 10'9, 11'5
τ Boötis . . . .	Apr.—June	13 41'8	+18 2	8'9	352'0	4'1, 11'5
94 Ceti . . . .	Dec.—Feb.	3 6'9	- 1 37	5'7	250'9	5'5, 11'5
κ Delphini . . . .	July—Nov.	20 33'6	+ 9 41	12'5	319'0	5'2, 11'8
α <sup>2</sup> Cancri . . . .	Jan.—Apr.	8 52'3	+12 18	11'1	325'0	4'4, 12'0
β Aquilæ . . . .	July—Oct.	19 49'7	+ 6 7	12'5	17'7	3'7, 12'1
B.A.C. 8173 . . . .	—	23 21'4	+70 3	20'1	312'3	6'5, 11'9
χ Delphini . . . .	July—Nov.	20 50'2	+12 7	40'0	21'8	6'0, 12'0
ξ Pegasi . . . .	Aug.—Nov.	22 41'0	+11 35	11'9, 127'3	112'6, 21'8	4'6, 12'2, 11'3
30 Pegasi . . . .	Aug.—Nov.	22 14'7	+ 5 13	6'3, 10'1	20'7, 222'8	6'0, 11'4, 12'3
53 Virginis . . . .	Apr.—June	13 6'0	-15 35	70'9	9'6	6'0, 12'5
30 Geminorum . . . .	Dec.—Apr.	6 37'5	+13 21	28'0	184'1	6'0, 12'7

## SELENOGRAPHICAL LONGITUDES AND LATITUDES OF LUNAR CRATERS.

NAME OF CRATER	Longitude		Latitude		NAME OF CRATER	Longitude		Latitude	
	+ = West	- = East	N = North	S = South		+ = West	- = East	N = North	S = South
Agrippa (102) i.	+ 10	22	N 4	4	Hesiodus (B) [187] iii.	- 17	0	S 26	50
Airy (291) iv.	+ 5	54	S 17	46	Hyginus (93) i.	+ 6	22	N 8	2
Albategnius (289) iv.	+ 3	58	S 11	21	Kepler (144) ii.	- 37	40	N 7	51
Alpetragius (205) iii.	- 4	45	S 15	55	La Hire (123) ii.	- 25	10	N 27	18
Alphonsus (207) iii.	- 3	14	S 12	59	Lalande (210) iii.	- 8	46	S 4	27
Archimedes (A) [120] ii.	- 7	11	N 27	45	Landsberg (222) iii.	- 26	20	S 0	27
Archytas (46) i.	+ 4	13	N 58	24	Langrenus (338) iv.	+ 60	30	S 8	33
Argelander (467) iv.	+ 41	43	S 45	46	Laplace (A) [134] ii.	- 26	34	N 43	16
Aristarchus (148) ii.	- 47	10	N 23	32	Le*Monnier (A) [53] i.	+ 29	8	N 26	0
Aristillus (83) i.	+ 1	1	N 33	45	Lichtenberg (151) ii.	- 67	5	N 31	25
Beer (446) iv.	+ 34	33	S 17	48	Lindenau (370) iv.	+ 24	30	S 31	52
Bessaron (145) ii.	- 37	4	N 14	52	Linné (74) i.	+ 11	33	N 27	48
Bessel (73) i.	+ 17	22	N 21	54	Mädler (466) iv.	+ 29	12	S 10	56
Billy (266) iii.	- 49	58	S 14	0	Manilius (95) i.	+ 8	47	N 14	27
Bode (107) ii.	- 2	39	N 6	37	Marius (147) ii.	- 50	6	N 11	50
Brayley (479) ii.	- 36	52	N 20	54	Maskelyne (67) i.	+ 29	40	N 2	32
Bullialdus (213) iii.	- 22	7	S 20	30	Menelaus (70) i.	+ 15	31	N 16	24
Burg (50) i.	+ 27	32	N 44	57	Messier (327) iv.	+ 47	12	S 1	57
Calippus (76) i.	+ 10	29	N 38	46	Milichius (118) ii.	- 29	40	N 10	0
Campanus (226) iii.	- 27	27	S 27	37	Moretus (262) iii.	- 7	9	S 69	45
Capella (324) iv.	+ 34	48	S 7	33	Mösting (211) iii.	- 5	54	S 0	36
Capuanus (238) iii.	- 25	42	S 34	20	Murchison (A) [483] i.	+ 1	0	N 4	6
Carlini (128) ii.	- 24	3	N 33	30	Mutus (400) iv.	+ 29	22	S 63	6
Cassini (A) [81] i.	+ 4	9	N 40	23	Neander (373) iv.	+ 39	45	S 31	10
Censorinus (325) iv.	+ 32	22	S 0	27	Olbers (159) ii.	- 77	33	N 7	55
Cepheus (31) i.	+ 45	40	N 40	59	Petavius (340) iv.	+ 59	47	S 24	50
Cleomedes (12) i.	+ 54	47	N 26	50	Picard (4) i.	+ 53	52	N 14	28
Conon (88) i.	+ 1	57	N 21	31	Piccolomini (371) iv.	+ 31	45	S 29	11
Copernicus (112) ii.	- 20	0	N 9	21	Pico (131) ii.	- 9	12	N 45	28
Crüger (278) iii.	- 66	40	S 16	48	Plinius (61) i.	+ 23	23	N 15	17
Delambre (301) iv.	+ 17	29	S 2	1	Posidonius (A) [54] i.	+ 29	11	N 31	33
Delisle (127) ii.	- 34	48	N 29	59	Proclus (60) i.	+ 46	30	N 16	10
Democritus (38) i.	+ 33	30	N 62	8	Prom. Agarum. (1) i.	+ 64	11	N 13	54
Dionysius (99) i.	+ 17	9	N 2	51	Pythagoras (176) ii.	- 62	15	N 63	6
Dollond (303) iv.	+ 14	16	S 10	14	Pytheas (124) ii.	- 20	30	N 20	14
Drebbel (240) iii.	- 48	13	S 40	47	Ramsden (228) iii.	- 31	42	S 32	26
Eichstädt (280) iii.	- 70	27	S 20	31	Reiner (146) ii.	- 56	1	N 7	5
Eimmart (3) i.	+ 62	50	N 23	35	Reinhold (114) ii.	- 22	37	N 3	13
Encke (143) ii.	- 36	37	N 4	18	Sacrobosco (312) iv.	+ 16	2	S 24	19
Eratosthenes (110) ii.	- 11	37	N 14	24	Scheiner (A) [261] iii.	- 26	36	S 59	58
Euclides (221) iii.	- 29	25	S 7	11	Schubert (A) [10] i.	+ 77	16	N 2	28
Euler (125) ii.	- 28	54	N 23	6	Seleucus (162) ii.	- 65	48	N 20	54
Fabricius (383) iv.	+ 40	46	S 42	8	Taruntius (326) i.	+ 45	59	N 5	40
Franklin (32) i.	+ 47	12	N 38	39	Thales (36) i.	+ 49	12	N 61	58
Flamsteed (223) iii.	- 44	12	S 4	31	Theophilus (319) iv.	+ 26	18	S 11	21
Gassendi (232) iii.	- 39	31	S 16	57	Timocharis (121) ii.	- 13	0	N 26	43
Goclenius (328) iv.	+ 44	29	S 10	0	Tobias Mayer (117) ii.	- 28	50	N 15	33
Grimaldi (272) iii.	- 68	58	S 2	43	Tycho (180) iii.	- 11	52	S 42	52
Hansen (A) [11] i.	+ 74	0	N 13	17	Ukert (109) i.	+ 1	9	N 7	48
Harding (152) ii.	- 70	52	N 43	9	Vitello (229) iii.	- 37	12	S 30	0
Helicon (129) ii.	- 22	53	N 40	10	Vitruvius (57) i.	+ 31	3	N 17	36
Hell (184) iii.	- 8	20	S 31	59	Werner (295) iv.	+ 3	6	S 27	30
Heraclides (135) ii.	- 31	1	N 41	8	Wichmann (438) iii.	- 37	56	S 7	41
Hercules (29) i.	+ 38	2	N 46	4	Wollaston (150) ii.	- 46	54	N 30	17
Herschel (212) iii.	- 2	9	S 5	37	Zagut (365) iv.	+ 21	56	S 31	42

1" of arc at the mean distance of the Moon is equal to 1.74 miles.

1° of selenographical longitude or latitude at the centre of the Moon is equal to 16"6 of arc, or 18.9 miles.

## LUNAR CRATERS IN ORDER OF SELENOGRAPHICAL LONGITUDE.

Longitude + = West	NAME OF CRATER	Latitude N = North S = South	Longitude - = East	NAME OF CRATER	Latitude N = North S = South
+77 16	Schubert (A) [10] i.	N 2 28	- 2 9	Herschel (212) iii.	S 5 37
74 0	Hansen (A) [11] i.	N 13 17	2 39	Bode (107) ii.	N 6 37
64 11	Prom. Agarum, (1) i.	N 13 54	3 14	Alphonsus (207) iii.	S 12 59
62 50	Eimmart (3) i.	N 23 35	4 45	Alpetragius (205) iii.	S 15 55
60 30	Langrenus (338) iv.	S 8 33	5 54	Mösting (211) iii.	S 0 36
59 47	Petavius (340) iv.	S 24 50	7 9	Moretus (262) iii.	S 69 45
54 47	Cleomedes (12) i.	N 26 50	7 11	Archimedes (A) [120] ii.	N 27 45
53 52	Picard (4) i.	N 14 28	8 20	Hell (184) iii.	S 31 59
49 12	Thales (36) i.	N 61 58	8 46	Lalande (210) iii.	S 4 27
47 12	Franklin (32) i.	N 38 39	9 12	Pico (131) ii.	N 45 28
47 12	Messier (327) iv.	S 1 57	11 37	Eratosthenes (110) ii.	N 14 24
46 30	Proclus (60) i.	N 16 10	11 52	Tycho (180) iii.	S 42 52
45 59	Taruntius (326) i.	N 5 40	13 0	Timocharis (121) ii.	N 26 43
45 40	Cepheus (31) i.	N 40 59	17 0	Hesiodus (B) [187] iii.	S 26 50
44 29	Goclenius (328) iv.	S 10 0	20 0	Copernicus (112) ii.	N 9 21
41 43	Argelander (467) iv.	S 45 46	20 30	Pytheas (124) ii.	N 20 14
40 46	Fabricius (383) iv.	S 42 8	22 7	Bullialdus (213) iii.	S 20 30
39 45	Neander (373) iv.	S 31 10	22 37	Reinhold (114) ii.	N 3 13
38 2	Hercules (29) i.	N 46 4	22 53	Helicon (129) ii.	N 40 10
34 48	Capella (324) iv.	S 7 33	24 3	Carlini (128) ii.	N 33 30
34 33	Beer (446) iv.	S 17 48	25 10	La Hire (123) ii.	N 27 18
33 30	Democritus (38) i.	N 62 8	25 42	Capuanus (238) iii.	S 34 20
32 22	Censorinus (325) iv.	S 0 27	26 20	Landsberg (222) iii.	S 0 27
31 45	Piccolomini (371) iv.	S 29 11	26 34	Laplace (A) [134] ii.	N 43 16
31 3	Vitruvius (57) i.	N 17 36	26 36	Scheiner (A) [261] iii.	S 59 58
29 40	Maskelyne (67) i.	N 2 32	27 27	Campanus (226) iii.	S 27 37
29 22	Mutus (400) iv.	S 63 6	28 50	Tobias Mayer (117) ii.	N 15 33
29 12	Mädler (466) iv.	S 10 56	28 54	Euler (125) ii.	N 23 6
29 11	Posidonius (A) [54] i.	N 31 33	29 25	Euclides (221) iii.	S 7 11
29 8	Le Monnier (A) [53] i.	N 26 0	29 40	Milichius (118) ii.	N 10 0
27 32	Burg (50) i.	N 44 57	31 1	Heraclides (135) ii.	N 41 8
26 18	Theophilus (319) iv.	S 11 21	31 42	Ramsden (228) iii.	S 32 26
24 30	Lindenau (370) iv.	S 31 52	34 48	Delisle (127) ii.	N 29 59
23 23	Plinius (61) i.	N 15 17	36 37	Encke (143) ii.	N 4 18
21 56	Zagut (369) iv.	S 31 42	36 52	Brayley (479) ii.	N 20 54
17 29	Delambre (301) iv.	S 2 1	37 4	Bessarion (145) ii.	N 14 52
17 22	Bessel (73) i.	N 21 54	37 12	Vitello (229) iii.	S 30 0
17 9	Dionysius (99) i.	N 2 51	37 40	Kepler (144) ii.	N 7 51
16 2	Sacrobosco (312) iv.	S 24 19	37 56	Wichmann (438) iii.	S 7 41
15 31	Menelaus (70) i.	N 16 24	39 31	Gassendi (232) iii.	S 16 57
14 16	Dollond (303) iv.	S 10 14	44 12	Flamsteed (223) iii.	S 4 31
11 33	Linné (74) i.	N 27 48	46 54	Wollaston (150) ii.	N 30 17
10 29	Calippus (76) i.	N 38 46	47 10	Aristarchus (148) ii.	N 23 32
10 22	Agrippa (102) i.	N 4 4	48 13	Drebbel (240) iii.	S 40 47
8 47	Manilius (95) i.	N 14 27	49 58	Billy (266) iii.	S 14 0
6 22	Hyginus (93) i.	N 8 2	50 6	Marius (147) ii.	N 11 50
5 54	Airy (291) iv.	S 17 46	56 1	Reiner (146) ii.	N 7 5
4 13	Archytas (46) i.	N 58 24	62 15	Pythagoras (176) ii.	N 63 6
4 9	Cassini (A) [81] i.	N 40 33	65 48	Seleucus (162) ii.	N 20 54
3 58	Albatagnius (289) iv.	S 11 21	66 40	Crüger (278) iii.	S 16 48
3 6	Werner (295) iv.	S 27 30	67 5	Lichtenberg (151) ii.	N 31 25
1 57	Conon (88) i.	N 21 31	68 58	Grimaldi (272) iii.	S 2 43
1 9	Ukert (109) i.	N 7 48	70 27	Eichstädt (280) iii.	S 20 31
1 1	Aristillus (83) i.	N 33 45	70 52	Harding (152) ii.	N 43 9
+ 1 0	Murchison (A) [483] i.	N 4 4	-77 33	Olbers (159) ii.	N 7 55

## SEPARATING POWER OF TELESCOPES.

Dawes (*Memoirs of the Royal Astronomical Society*, volume xxxv. p. 158) found by a great variety of experiments with small telescopes that a one-inch aperture would just separate a double star composed of two stars of the sixth magnitude if their central distance was  $4''\cdot56$ , when the atmospheric circumstances were moderately favourable. The following little table, as calculated by him, will be convenient for reference.

Aperture in Inches	Least separable Distance	Aperture in Inches	Least separable Distance	Aperture in Inches	Least separable Distance
1'0	$4''\cdot56$	4'0	$1''\cdot14$	7'0	$0''\cdot65$
1'5	$3''\cdot04$	4'5	$1''\cdot01$	8'0	$0''\cdot57$
2'0	$2''\cdot28$	5'0	$0''\cdot91$	9'0	$0''\cdot507$
2'5	$1''\cdot82$	5'5	$0''\cdot83$	10'0	$0''\cdot456$
3'0	$1''\cdot52$	6'0	$0''\cdot76$	11'0	$0''\cdot414$
3'5	$1''\cdot30$	6'5	$0''\cdot70$	12'0	$0''\cdot380$

## SPACE-PENETRATING POWER OF TELESCOPES.

The following table, based on the assumption that a star of any magnitude contains  $2\frac{1}{2}$  times the light of one of the magnitude next below, is taken from Newcomb and Holden's *Astronomy*, 4th edition, page 419. This is practically Struve's ratio.

Aperture in Inches	Faintest Star Visible	Aperture in Inches	Faintest Star Visible	Aperture in Inches	Faintest Star Visible
1'0	Mag. 9'0	4'0	Mag. 12'0	7'0	Mag. 13'3
1'5	9'9	4'5	12'3	8'0	13'5
2'0	10'5	5'0	12'5	9'0	13'8
2'5	11'0	5'5	12'7	10'0	14'0
3'0	11'4	6'0	12'9	11'0	14'2
3'5	11'7	6'5	13'1	12'0	14'4

## APPENDIX.

Heliocentric Minima of  $\beta$  Persei (*Algol*)  $3^h 0^m 45^s + 40^\circ 31' 0''$ .

	Days hrs. min.	Days hrs. min.	Days hrs. min.	Days hrs. min.
1886, Jan.	13 14 23	March 6 5 3	August 16 15 30	October 24 11 4
	16 11 12	20 13 7	19 12 19	27 7 53
	19 8 1	23 9 56	22 9 8	November 13 12 46
	22 4 50	April 9 14 50	September 8 14 1	16 9 35
February	5 12 55	12 11 39	11 10 50	19 6 24
	8 9 43	15 8 27	14 7 39	December 3 14 28
	11 6 32	July 7 12 6	October 1 12 32	6 11 17
	25 14 36	10 8 55	4 9 21	9 8 6
	28 11 25	27 13 29	7 6 10	26 12 59
March	3 8 14	30 10 37	21 14 15	29 9 48

*Algol*, 2.2 mag. at maximum, 3.7 at minimum. It commences to diminish in light 4 hours 23 minutes before minimum, and regains its full brilliancy 5 hours and 37 minutes after. The most rapid decrease of light begins at 1 hour 40 minutes before the minimum, and the most rapid increase 1 hour 40 minutes after.

Heliocentric Minima of  $\delta$  Libræ  $14^h 54^m 53^s - 8^\circ 3' 9''$ . 4.9 mag. max. 6.1 mag. min.

	Days hrs. min.	Days hrs. min.	Days hrs. min.	Days hrs. min.
April	3 16 44	May 1 15 0	June 5 12 50	July 3 11 6
	10 16 18	8 14 34	12 12 24	10 10 40
	17 15 52	15 14 8	19 11 58	17 10 14
	24 15 26	22 13 42	26 11 32	24 9 48
		29 13 16		31 9 22



# VARIABLE STARS.

NAME OF VARIABLE STAR.	R.A.		Decl.	MAGNITUDE		EPHEMERIS (1886)		NOTES AND COLOURS
	h. m. s.			Maximum	Minimum	Maximum	Minimum	
	R. Andromedæ . . . . .	0		18	1	5'6-8'6	Below 13	
R. Arietis . . . . .	2	9	38	7'6-8'6	12.3	February 6, August 11	{ November 10 (1885), May 15, Nov. 17 }	Close to the 6th mag. star 21 Arietis.
o Ceti ( <i>Mira</i> ) . . . . .	2	13	35	1'7-5	8-9.5	January 5	August 15	Very red near minimum, a blue 9.5 mag. at 8 <sup>o</sup> : 116'.
R. Leporis . . . . .	4	54	25	6.7	8.5	December 12	April 26	Hind's scarlet star. Irregular. In 1883 the maxima and minima took place 8 weeks before the predicted times.
S. Orionis . . . . .	5	23	23	8.5	Below 13	November 1	April 15	Very red. Period rather uncertain.
R. Leonis . . . . .	9	41	26	5'2-6'4	9.4-10	{ November 9 (1885), September 18 }	April 23	Very red.
R. Ursæ Maj. . . . .	10	36	34	6'1-8'1	13.4	May 11	January 28	Orange.
R. Virginis . . . . .	12	32	43	6'5-7'5	10-10.9	{ November 15 (1885), April 9, September 2 }	{ January 31, June 25, November 18 }	Reddish at maximum, deeper at minimum, like many other variables.
R. Hydræ . . . . .	13	23	30	4-5.5	10.5	November 2	April 16	Red. P. xiii. 94.
R. Aquilæ . . . . .	19	0	53	6'4-7'4	11.2	May 11	November 21	Reddish.
* Cygni . . . . .	19	46	11	4'0-6'0	12.8	January 8	July 19	x of Bayer, not x of Flamsteed. Fine red.
T. Cephei . . . . .	21	8	2	6.4	9.8	March 13	September 20	Reddish.





## THE COMETS OF 1886.

Three periodical comets are expected to return to perihelion in the year 1886. These are (1) a comet having a period of about  $5\frac{1}{2}$  years, which was discovered by Tempel at Florence in November 1869. The return in 1875 was not observed, but it was re-discovered by Prof. L. Swift, of Rochester, New York, in October 1880, when the fact of its periodicity was determined. Perihelion passage will take place about April 21, 1886, but its position in the heavens will be very unfavourable for observation, and it is not likely to be seen again until its next return in October 1891.

(2) A comet, first discovered by Pons at Marseilles in June 1819, which is also moving in an ellipse with a period of about  $5\frac{1}{2}$  years. It was re-discovered by Dr. Winnecke in 1858, and its periodicity was then determined by him. It was well observed in 1869 and 1875, but not in 1864 and 1880, when the circumstances were unfavourable. It is usually known as Winnecke's comet. Perihelion passage will take place about August 30, 1886.


(3) A comet of long period, discovered by Olbers in March 1815, is expected to return again to perihelion towards the end of 1886. It will probably be bright enough to be generally observed. A sweeping ephemeris from January to October 1886 has been published by Dr. Ginzell in No. 2,696 of the *Astronomische Nachrichten*.

Three comets, discovered in 1885, are visible at the present time. They are :—

(1) A comet discovered by Mr. Brooks, of Phelps, New York, at the end of December 1885. This comet passed its perihelion on November 24, 1885, and is slowly receding from the earth, and growing fainter.

(2) A comet discovered by M. Fabry at the Paris Observatory on December 1, 1885. This comet is only observable with large telescopes. Perihelion passage, according to the elements calculated by Dr. Oppenheim, will take place on April 4, 1886. The brightness is slowly increasing at present.

(3) A comet discovered by Mr. Barnard, at the Vanderbilt University Observatory, Nashville, Tennessee, on December 3, 1885. This comet is also very faint. Perihelion passage will take place about May 5, 1886. The elements bear some resemblance to those of the second comet of 1785. Its brightness is slowly increasing at present.




## ADDENDA.

THE orbit of comet Fabry has been recently calculated by several astronomers, including Dr. S. Oppenheim and M. Lebeuf, and it appears from their investigations that this comet, instead of decreasing in brightness, promises to be a very conspicuous object in the circumpolar sky towards the end of April and the beginning of May. According to Dr. Oppenheim its brightness on May 1 will be over six hundred times greater than that at the time of discovery.

The orbit of comet Barnard has also been investigated by Dr. Heppenger, with a somewhat similar result. About the middle of May this comet will be nearly three hundred times brighter than at the date of discovery.



# BROWNING'S ASTRONOMICAL TELESCOPES.



## ASTRONOMICAL TELESCOPE

With 3-inch Achromatic Object-Glass of excellent quality, with brass body, one Terrestrial and two Celestial Eyepieces, in case complete. Guaranteed to be capable of dividing Double-Stars and showing Saturn's Ring and Jupiter's Belts.

Price £6. 10s.

*Catalogue of Astronomical Telescopes sent Free.*

## SILVERED GLASS TELESCOPES AND SPECULA.

### SILVERED GLASS SPECULA, UNMOUNTED.

#### WITHOUT CELLS.

To meet the wishes of many applicants Mr. John Browning has made arrangements with Mr. With whereby the large stock of Specula, now in the possession of that celebrated artist, may be obtained by those who wish to possess one of Mr. With's Specula, but to do the mounting for themselves. The Specula so available are the choicest reserves and re-touches of Mr. With's productions.

To assist amateurs to the very utmost Mr. Browning will supply warranted planes, finders, eye-pieces, &c., of his own manufacture, indeed all those parts of the instrument which an amateur cannot as a whole make for himself.

The performance of these Specula will be guaranteed; they will bear a power of 100 to the inch on suitable objects, and under favourable conditions of the atmosphere.

Speculum, 4½-inch diameter about 5 ft. focus, £2. 10s.; 6½-inch, about 6 ft. focus, £5; 8½-inch, £8; 9½-inch, £12; 10½-inch, £16; 12½-inch, £22; 13-inch, £25.

### SILVERED GLASS SPECULA ASTRONOMICAL TELESCOPES, ON ALT-AZIMUTH STANDS.

4½-inch Speculum, 5 ft. focus, mounted on a stand, which can be changed from alt-azimuth to parallaxic, so that the stars can be followed with one motion, with endless driving screw, and Hook's joint and two eye-pieces, 100 to 200 ... ..	£	s.	d.
6½-inch Speculum, 6 ft. focus, on alt-azimuth stand, with quick and slow fine screw motions, and three eye-pieces, 100 to 450 ... ..	20	0	0
	31	0	0

*Full list of Prices sent free.*

#### EXTRACTS FROM

'A PLEA FOR REFLECTORS,' being a description of the new Astronomical Telescopes, with silvered glass specula, and instructions for using and adjusting them by JOHN BROWNING, F.R.A.S.—Post free.

JOHN BROWNING, 63 Strand, London, W.C.

# J. H. DALLMEYER,

19 BLOOMSBURY STREET, LONDON, W.C.

## INTERNATIONAL EXHIBITION, 1862.

“The Highest Award for his excellent Object Glasses and Equatorial Mountings.”

## PARIS INTERNATIONAL EXHIBITION, 1867.

“The Gold Medal for the best Telescope, and the Silver Medal for the best Photographic Lens.”

---

### REPORT BY ORDER OF THE COUNCIL ON EDUCATION.

“It is satisfactory to be able to record that the result of a trial by the jurors of the Refracting Astronomical Telescopes is to place Mr. Dallmeyer at the head of the list. The performance, power, and definition of his Equatorial, surpasses that of the other astronomical telescopes.”

“J. H. Dallmeyer has been very successful in producing hand telescopes of a superior description; he exhibits some instruments, the focal lengths of which are only from nine to ten times the diameter of their clear aperture. The performance of these instruments is most excellent.”—ILLUSTRATED LONDON NEWS, Oct. 5, 1867 (page 378).

---

## PHILADELPHIA EXHIBITION, 1876.

“Highest Award for Telescopes, Microscopes, Photographic Lenses, and Apparatus.”

## PARIS INTERNATIONAL EXHIBITION, 1878.

“The Cross of the Legion of Honour and Two Gold Medals.” Exhibit: Telescopes, Microscopes, Photographic Lenses, &c.

---

# PHOTOHELIOGRAPHS.

*Supplied to the Home and Foreign Governments.*

---

## CENTENNIAL EXHIBITION, PHILADELPHIA, 1876.

“Some very beautiful sharply defined pictures of the Sun, with interesting groups of Sun spots, are shown, and furnish a very satisfactory proof of the excellent Photoheliographic Apparatus, by means of which they were obtained.”—*Vide* JUROR'S REPORT.

---

TRANSIT INSTRUMENTS, STAR-SPECTROSCOPES, &c.  
*Instruments for Stellar Photography.*

---

*Descriptive Catalogue of Telescopes, Microscopes, Photographic Lenses, &c., on application.*

---

# J. H. DALLMEYER,

19 BLOOMSBURY STREET, LONDON, W.C.




A. J. FROST, 6 Westminster Chambers, Victoria Street, S.W.

# CLARK'S PATENT IMPROVED TRANSIT INSTRUMENTS.

*International Inventions Exhibition, 1885, Medal awarded.*

13 inch, with striding level, lamp, and support,  
complete £9. 17s. 6d.



18 inch, with striding level, axial illumination, and lamp  
and support, complete £14. 14s. 0d.


These Instruments are of high quality, accurately adjusted, and well finished. The base is of cast iron, with large gun-metal levelling screws. The axis is of gun-metal, the trunnions resting on gun-metal bearings; the altitude or declination circle is silver-plated, and divided into 4 quadrants of 90 degrees with a vernier and spirit level reading to single minutes. It is furnished with a diagonal eye-piece with total reflecting prism, and three spider-web cross-wires, with collimating screws. There are two dark glasses for solar observation, and a delicate striding level is supplied with the Instrument. The optical power of the larger Instrument is sufficient to show the larger stars and planets in full daylight. The illumination of the cross-wires at night is effected by a front reflecting diaphragm, and the Instrument is provided with a candle lantern carried on an adjustable support. The ordinary axial illumination is supplied with the larger Instrument in addition.

## CLARK'S WINDOW TRANSIT.

In addition to the above instrument another form can be supplied for the use of amateurs desiring a small portable and easily-adjusted Instrument. This Instrument has been designed for fixing upon a window-sill and requires no special stand. The base is entirely of gun-metal, and is handsomely finished, Price £7. 10s. 0d.

*Pamphlet post-free on application.*

A. J. FROST, 6 Westminster Chambers, London, S.W.



## MAWSON & SWAN'S ASTRONOMICAL TELESCOPE.

Complete, with Stand and Box, £5.

Of superior construction. Is fitted with  $2\frac{3}{4}$ -inch Object Glass of Best English manufacture. Day and Night Eye-pieces. Highly finished Brass Stand, &c. Complete in box.

A cheaper form, fitted with Day and Night Eye-pieces, 2-inch Object Glass of good quality, French make, price £2. 15s.

*A very useful Instrument.*

---

## SWAN LAMPS FOR ASTRONOMICAL TELESCOPES.

The small  $2\frac{1}{2}$ -candle Swan Lamp has been most successfully employed for lighting up the Micrometer Scale &c., in Telescopes in many of the large and most important Observatories.

The four or five-cell **modified** Leclanché Battery, controlled by a simple switch, answers admirably for this work.

---

## PORTABLE PRIMARY BATTERIES,

*For working the small Swan Electric Lamps.*

---

*Catalogues of Chemical, Photographic, Electrical and Microscopical Apparatus on application.*


---

MAWSON & SWAN, Mosley Street, Newcastle-on-Tyne.

# THE 'COLLEGE' TELESCOPE.

This Instrument has a brass body, with rack and pinion adjustment ; is 38-in. long and 3-in. in diameter. Mounted on a brass pillar, with horizontal and vertical movements and japanned claw stand. The Day Power is 40-dia., and the Huyghenian Astro Eyepiece magnifies 90-dia. The latter is fitted with a dark glass.

Price



£5. 5s.

**PACKED IN PORTABLE PINE CASE.**

ILLUSTRATED CATALOGUE OF ASTRONOMICAL TELESCOPES, JUST PUBLISHED; GRATIS, ON APPLICATION, OR BY POST ONE STAMP.

**HORNE, THORNTHWAITE, & WOOD,**  
416 STRAND, W.C., AND  
**E. G. WOOD,**  
74 CHEAPSIDE, E.C.

## W. WRAY, Optician,

MANUFACTURER OF

### ASTRONOMICAL TELESCOPES.

*EQUATORIAL and other MOUNTINGS, EYEPIECES, &c. &c.*

MICROSCOPE OBJECTIVES AND MICROSCOPES.

PHOTOGRAPHIC LENSES, &c. &c.

Illustrated Catalogue Free.

LAUREL HOUSE, NORTH HILL, HIGHGATE,  
LONDON, N.

# WORKS ON THE TRANSIT INSTRUMENT.

By LATIMER CLARK, M.I.C.E.

---

Crown 8vo. price 1s.

## MANUAL OF THE TRANSIT INSTRUMENT.

This work gives all the information requisite for taking time by the Transit Instrument. By the aid of the Tables mentioned below, anyone can take transits *without any previous knowledge of the subject.*

---

Demy 8vo. cloth 5s.

## A TREATISE ON THE TRANSIT INSTRUMENT

*As applied to the determination of Time.*

This work gives a complete description of the Transit Instrument, and the adjustments and corrections required for obtaining accurate time in any part of the world. It forms an excellent practical introduction to the science of astronomy.

---

Crown 8vo. boards, price 2s. 6d. annually.

## TRANSIT TABLES.

These Tables, which are published annually, give the transit of about twenty selected stars for every evening in the year in ordinary time, so that in taking time by the Transit Instrument no calculation of any kind is required. They are adapted for use in all parts of the globe.

---

LONDON: E. & F. N. SPON, 125 Strand.

NEW YORK: 35 Murray Street







14 DAY USE  
RETURN TO DESK FROM WHICH BORROWED

**LOAN DEPT.**

This book is due on the last date stamped below, or  
on the date to which renewed.

Renewed books are subject to immediate recall.

9 FEB '60 AR

REC'D LD

JAN 27 1960

DEC 31 1968 25

RECEIVED

DEC 20 '68 -1 PM

LOAN DEPT.

LD 21A-50m-4,'59  
(A1724s10)476B

General Library  
University of California  
Berkeley



U. C. BERKELEY LIBRARY



061357053

QB64  
C6

127067

Clark

